



ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS

ACTIVE CONSTITUENTS: Each tablet contains 545mg OXANTEL EMBONATE, 140mg PYRANTEL EMBONATE and 50mg PRAZIQUANTEL

Cost effective broad spectrum wormer for dogs

For treatment and control of roundworms (*Toxocara spp.*, *Toxascaris leonina*), hookworms (*Uncinaria stenocephala*, *Ancylostoma spp.*), whipworms (*Trichuris vulpis*) and adult tapeworms (*Taenia spp.*, *Dipylidium caninum*, *Spirometra erinacei* and *Echinococcus granulosus*) in dogs.

NEW
BLISTER
PACK

- 🐾 Broad spectrum intestinal wormer
- 🐾 Easy to administer
- 🐾 Scored tablets for targeted dosing
- 🐾 Cost effective
- 🐾 Suitable for all dog sizes
- 🐾 Safe for puppies from 2 weeks
- 🐾 New blister pack design offering:
 - Enhanced individual tablet protection
 - Reduced risk of breakage
 - Improved tamper-evidence



	ABANTEL™	Popantel®
Species	Dog	Dog
Route of Administration	Tablet	Tablet
Tablet dose	1 tablet per 10kg bodyweight dog	1 tablet per 10kg bodyweight dog
Active Constituents		
Praziquantel	50mg	50mg
Pyrantel Embonate	140mg	143mg
Oxantel Embonate	545mg	542mg
Controls Roundworms, Whipworms, Hookworms & Tapeworms	Yes	Yes

ABANTEL™ is a trademark of Abbey Laboratories Pty Ltd

Popantel® is a registered trademark of Zoetis Services LLC

ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS

ACTIVE CONSTITUENTS: Each tablet contains 545mg OXANTEL EMBONATE, 140mg PYRANTEL EMBONATE and 50mg PRAZIQUANTEL



For treatment and control of roundworms (*Toxocara spp.*, *Toxascaris leonina*), hookworms (*Uncinaria stenocephala*, *Ancylostoma spp.*), whipworms (*Trichuris vulpis*) and adult tapeworms (*Taenia spp.*, *Dipylidium caninum*, *Spirometra erinacei* and *Echinococcus granulosus*) in dogs.

Pyrantel embonate and oxantel embonate belong to the tetrahydropyrimidine group. Praziquantel is a pyrazion-isoquinolein derivative belonging to the thioxanonic group.

This product does not control heartworm.

If worm problems persist, consult a veterinarian.

For use by or under direction of a veterinarian.

DIRECTIONS FOR USE

DOSAGE AND ADMINISTRATION

For oral use only. Can be administered directly into the mouth or crushed and mixed with the feed.

1 tablet per 10kg bodyweight.

Dogs up to 5kg should be given half (½) a tablet.

Dosage table:

Bodyweight	Dose
up to 5kg	½ tablet
10kg	1 tablet
20kg	2 tablets
30kg	3 tablets
40kg	4 tablets
50kg	5 tablets

Specific treatment for *Spirometra erinacei* is by administering ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS four (4) times the above dose rate.

GENERAL DIRECTIONS

ROUNDWORMS & HOOKWORMS:

Treat at 2, 4, 8 and 12 weeks - thereafter every 3 months.

WHIPWORMS: Treat every 6-8 weeks after 3 months of age.

HYDATID TAPEWORM: Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks.

OTHER TAPEWORMS: Treat every 3 months.

PREGNANT BITCHES: Treat at mating, before whelping and then every 3 months.

For *Dipylidium* treatment adequate flea control should be maintained all year around.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

DISPOSAL

Dispose of empty container by wrapping with paper and placing in garbage.

STORAGE

Store below 25 °C (Air conditioning) in a dry place.

**APVMA Approval Number:
69893/149063**

