



# ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS

ACTIVE CONSTITUENTS: Each tablet contains - 545mg OXANTEL EMBONATE, 140mg PYRANTEL EMBONATE and 50mg PRAZIQUANTEL

## Cost effective broad spectrum wormer for dogs

For treatment and control of roundworms (*Toxocara spp.*, *Toxascaris leonina*), hookworms (*Uncinaria stenocephala*, *Ancylostoma spp*), whipworms (*Trichuris vulpis*) and adult tapeworms (*Taenia spp*, *Dipylidium caninum*, *Spirometra erinacei* and *Echinococcus granulosus*) in dogs.

- 🌿 Broad spectrum intestinal wormer
- 🌿 Easy to administer
- 🌿 Scored tablets for targeted dosing
- 🌿 Cost effective



	ABANTEL™	Popantel®
Species	Dog	Dog
Route of Administration	Tablet	Tablet
Tablet dose	1 tablet per 10kg bodyweight dog	1 tablet per 10kg bodyweight dog
Active Constituents		
Praziquantel	50mg	50mg
Pyrantel Embonate	140mg	143mg
Oxantel Embonate	545mg	542mg
Controls Roundworms, Whipworms, Hookworms & Tapeworms	Yes	Yes

ABANTEL™ is a trademark of Abbey Laboratories Pty Ltd

Popantel® is a registered trademark of Jurox Pty Ltd

# ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS

ACTIVE CONSTITUENTS: Each tablet contains - 545mg OXANTEL EMBONATE, 140mg PYRANTEL EMBONATE and 50mg PRAZIQUANTEL



**For treatment and control of roundworms (*Toxocara spp.*, *Toxascaris leonina*), hookworms (*Uncinaria stenocephala*, *Ancylostoma spp*), whipworms (*Trichuris vulpis*) and adult tapeworms (*Taenia spp*, *Dipylidium caninum*, *Spirometra erinacei* and *Echinococcus granulosus*) in dogs.**

This product does not control heartworm.

If worm problems persist, consult a veterinarian.

## DIRECTIONS FOR USE

### DOSAGE AND ADMINISTRATION

For oral use only. Can be administered directly into the mouth or crushed and mixed with the feed.

1 tablet per 10kg bodyweight.

Dogs up to 5kg should be given half (½) a tablet.

### Dosage table:

Bodyweight	Dose
up to 5kg	½ tablet
10kg	1 tablet
20kg	2 tablets
30kg	3 tablets
40kg	4 tablets
50kg	5 tablets

Specific treatment for *Spirometra erinacei* is by administering ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS four (4) times the above dose rate.

## GENERAL DIRECTIONS

### ROUNDWORMS & HOOKWORMS:

Treat at 2, 4, 8 and 12 weeks - thereafter every 3 months.

**WHIPWORMS:** Treat every 6-8 weeks after 3 months of age.

**HYDATID TAPEWORM:** Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks.

**OTHER TAPEWORMS:** Treat every 3 months.

**PREGNANT BITCHES:** Treat at mating, before whelping and then every 3 months.

For *Dipylidium* treatment adequate flea control should be maintained all year around.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 13 11 26.

## DISPOSAL

Dispose of empty container by wrapping with paper and placing in garbage.

## STORAGE

Store below 25 °C (Air conditioning) in a dry place.

**APVMA Approval Number:  
69893/111735**