








ABBEY™ TIAMULIN PREMIX

ACTIVE CONSTITUENT: 25g/kg TIAMULIN HYDROGEN FUMARATE

Targeted treatment for swine and poultry

For the treatment of swine dysentery caused by *Brachyspira hyodysenteriae*, and Swine Enzootic Pneumonia caused by *Mycoplasma hyopneumoniae* and as an aid for the control of Swine Enzootic Pneumonia caused by *Mycoplasma hyopneumoniae* in growing pigs. For the treatment of Air Sacculitis (Chronic Respiratory Disease) caused by *Mycoplasma gallisepticum* in non-laying chickens.

-  Easy to use – just mix in feed
-  Dual species registration
-  Targeted treatment for bacterial and mycoplasma infections
-  Cost effective



ABBEY™ TIAMULIN PREMIX

Active Constituent	25g/kg TIAMULIN HYDROGEN FUMARATE
Species	Swine and Non-Laying Poultry
Route of Administration	Oral Premix
Dose Rate	
Pigs: Treatment of Swine Dysentery	8kg per 1000kg feed continuously for 5 days
Pigs: Treatment of Swine Enzootic Pneumonia	8kg per 1000kg feed continuously for 10 days
Pigs: Aid in control of Swine Enzootic Pneumonia	1kg per 1000kg feed
Non-Laying Poultry: Treatment Air Sacculitis (Chronic Respiratory Disease)	8kg per 1000kg feed continuously for 5 days
Withholding Period	
Meat	5 days
Eggs	Do not use

ABBEY™ is a trademark of Abbey Laboratories Pty Ltd

ABBEY™ TIAMULIN PREMIX

ACTIVE CONSTITUENT: 25g/kg TIAMULIN HYDROGEN FUMARATE



For the treatment of swine dysentery caused by *Brachyspira hyodysenteriae*, and Swine Enzootic Pneumonia caused by *Mycoplasma hyopneumoniae* and as an aid for the control of Swine Enzootic Pneumonia caused by *Mycoplasma hyopneumoniae* in growing pigs. For the treatment of Air Sacculitis (Chronic Respiratory Disease) caused by *Mycoplasma gallisepticum* in non-laying chickens.

DIRECTIONS FOR USE

Restraints

DO NOT USE in birds which are producing or may in the future produce eggs or egg products for human consumption.

Contraindications

This product is contraindicated for use in livestock receiving products containing Narasin, Monensin, or Salinomycin in the feed. During treatment withhold all other sources of feed. Tiamulin should not be given for 5-7 days of Narasin, Monensin or Salinomycin use.

DOSAGE AND ADMINISTRATION

ABBEY™ TIAMULIN PREMIX MUST BE THOROUGHLY MIXED IN FEED BEFORE USE.

PIGS	INCLUSION RATE
Treatment of clinical Swine Dysentery	8kg of ABBEY™ Tiamulin Premix per 1000kg feed. To be administered continuously for 5 days
Treatment of Swine Enzootic Pneumonia	8kg of ABBEY™ Tiamulin Premix per 1000kg feed. To be administered continuously for 10 days.
Aid in control of Swine Enzootic Pneumonia	1kg ABBEY™ Tiamulin Premix per 1000kg feed.
NON-LAYING POULTRY	INCLUSION RATE
Treatment Air Sacculitis (Chronic Respiratory Disease)	8kg of ABBEY™ Tiamulin Premix per 1000kg feed. To be administered continuously for 5 days.

GENERAL DIRECTIONS

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 5 days before slaughter for human consumption.

EGGS: DO NOT USE in birds which are producing or may in the future produce eggs or egg products for human consumption.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on ESI, contact Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website before using this product.

SAFETY DIRECTIONS

May irritate the eyes, avoid contact with eyes and skin. Do not inhale DUST when opening the container or mixing. Wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and disposable dust mask if dust is present. Wear goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

DISPOSAL

Shake empty bag into medicated feed. Shred and bury bags in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit. DO NOT burn empty containers or product.

STORAGE

Store below 30° C (Room Temperature) in a dry place.

APVMA Approval Number: 62394/105777