







ABBEYMOX™ FG Feed Grade Amoxicillin Soluble Powder

ACTIVE CONSTITUENT: 870g/kg AMOXYCILLIN AS THE TRIHYDRATE

Water Soluble Microbial

AbbeyMox™ FG Feed Grade Amoxicillin Soluble Powder is a broad spectrum water-soluble medication for the treatment of Amoxicillin susceptible infections of the alimentary, urogenital and respiratory tract of poultry, including organisms associated with Chronic Respiratory Disease (CRD) complex and diseases caused by *E. Coli*.

-  East to use - just add water!
-  Broad spectrum antibiotic
-  Treats a variety of systemic infection in poultry
-  Cost Effective



ABBEYMOX™ FG

Active Constituent	870g/kg Amoxicillin as the Trihydrate
Species	Poultry
Dose rate	8-20 mg/kg depending on severity of condition
Administration method	Soluble Powder
Withholding Period	
Meat	2 days
Eggs	Zero days

ABBEYMOX™ is a trademark of Abbey Laboratories Pty Ltd

ABBEYMOX™ FG Feed Grade Amoxicillin Soluble Powder

ACTIVE CONSTITUENT: 870g/kg AMOXYCILLIN AS THE TRIHYDRATE



A broad spectrum water-soluble medication for the treatment of Amoxicillin susceptible infections of the alimentary, urogenital and respiratory tract of poultry, including organisms associated with Chronic Respiratory Disease (CRD) complex and diseases caused by *E. Coli*.

DOSAGE AND ADMINISTRATION

ABBEYMOX™ FG Feed Grade Amoxicillin Soluble Powder is for use as a water medication for poultry species. Water consumption varies with the climate and the age of the birds. General dose rate of Amoxicillin for poultry is 8-20mg/kg bodyweight daily, depending on the severity of the condition. This may be achieved by adding 110g of ABBEYMOX™ FG Feed Grade Amoxicillin Soluble Powder to 900-1000L of drinking water for 3-5 days.

The material dissolves and disperses readily in water. Administer as the only source of drinking water for 3-5 days. Medicated water should be consumed within 12 hours, and fresh solution prepared daily. Water tanks and lines should be flushed thoroughly to avoid any holdover of residues of the product.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 2 days before slaughter for human consumption.

CHICKEN EGGS: Zero (0) days

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

TRADE ADVICE

Export Egg Withholding Interval: 14 days.

DO NOT ALLOW eggs treated with ABBEYMOX™ FG Feed Grade Amoxicillin Soluble Powder within the last 14 days to be sold for export.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. If product gets into eyes, wash it out immediately with water. Repeated exposure may cause allergic disorders.

Do not inhale dust and wash hands after use when opening the container and mixing into water. Wear cotton overalls buttoned to the neck and wrist, elbow-length PVC gloves, goggles and disposable dust face mask covering mouth and nose. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre: Phone Australia 131 126

DISPOSAL

Single rinse or shake remainder into the medicated water.

Do not dispose of undiluted chemicals on-site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature).

Store in a tightly closed container in a dry place.

APVMA Approval Number: 68888 / 141148