







# ABBEYMOX™ Amoxicillin Soluble Powder

ACTIVE CONSTITUENT: 870g/kg AMOXYCILLIN (as the Trihydrate)

## Water Soluble Microbial

Water soluble medication for the treatment of amoxicillin susceptible infections of the alimentary, urogenital and respiratory tracts of poultry, including organisms associated with the Chronic Respiratory Disease (CRD) complex and diseases caused by *E. Coli*.

-  East to use - just add to drinking water!
-  Broad spectrum antibiotic
-  Zero day egg WHP
-  Cost Effective



	ABBEYMOX™ AMOXYCILLIN SOLUBLE POWDER	CCD® AMOXYCILLIN TRIHYDRATE FOR POULTRY
Active Constituent	870g/kg Amoxicillin (as the Trihydrate)	870mg/g Amoxicillin (as Amoxicillin Trihydrate)
Species	Poultry	Poultry
Dose rate	8-20 mg/kg	8-20 mg/kg
Administration method	Water soluble	Water soluble
Withholding Period		
Meat	2 days	2 days
Eggs	Zero days	Zero days

ABBEYMOX™ is a trademark of Abbey Laboratories Pty Ltd

CCD® is a registered trademark of CCD Animal Health Pty Ltd

# ABBEYMOX™ Amoxicillin Soluble Powder

ACTIVE CONSTITUENT: 870g/kg AMOXYCILLIN (as the Trihydrate)



**Water soluble medication for the treatment of amoxicillin susceptible infections of the alimentary, urogenital and respiratory tracts of poultry, including organisms associated with the Chronic Respiratory Disease (CRD) complex and diseases caused by *E. Coli*.**

## DIRECTIONS FOR USE

### DOSAGE AND ADMINISTRATION

Dose for chickens, turkeys and other poultry: 8 - 20 mg Amoxicillin per kilogram live weight daily depending on the severity of the condition. This is achieved by adding 110 g of ABBEYMOX™ Amoxicillin Antibiotic Soluble Powder to 900-1000 litres of drinking water for 3 to 5 days. Medicated water should be consumed within 12 hours and fresh solution prepared daily. Remove other sources of medicated water.

## WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 2 days before slaughter for human consumption.**

**EGGS: Zero (0) days**

**Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.**

## TRADE ADVICE

**Export Egg Withholding Interval: 14 days.**

**DO NOT ALLOW eggs treated with ABBEYMOX™ Amoxicillin Antibiotic Soluble Powder within the last 14 days to be sold for export.**

## SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. If product gets into eyes, wash it out immediately with water. Repeated exposure may cause allergic disorders.

Do not inhale dust and wash hands after use when opening the container and mixing into water.

Wear cotton overalls buttoned to the neck and wrist, elbow-length PVC gloves, goggles and disposable dust face mask covering mouth and nose. After each day's use, wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre: Phone Australia 131 126.

## DISPOSAL

Single-rinse or shake remainder into the medicated water. Do not dispose of undiluted chemicals on-site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## STORAGE

Store in a dry place below 25 °C (air conditioning). Protect from light. Keep bag tightly closed.

**APVMA Approval Number: 93117/140579**