








AbbeyNeo™ Antibiotic Feed Additive

ACTIVE CONSTITUENT: 600g/kg NEOMYCIN (as Sulfate)

The easy option for Antibiotics

Broad-spectrum aminoglycoside antibiotic for the treatment of bacterial scours caused by organisms sensitive to Neomycin, including Neomycin sensitive strains of *E. coli*, *Salmonella* spp., *Pseudomonas* spp., in poultry, pigs, and cattle.

-  East to use - just mix into feed!
-  Suppresses growth of harmful bacteria in intestines
-  Reduce the incidence of infectious diseases
-  NIL milk and egg WHP



AbbeyNeo™	
Active Constituent	600g/kg Neomycin (as sulfate)
Species	Poultry, Pigs, Cattle
Dose rate	8-22 mg/kg for 3-5 days
Route of Administration	Feed Additive
Pack size	1kg

Withholding Periods	
Meat: Broilers	5 days
Meat: Turkeys	14 days
Meat: Laying Hens	14 days
Meat: Pigs	20 days
Meat: Cattle	30 days
Milk	0 days
Eggs	0 days

AbbeyNeo™ is a trademark of Abbey Laboratories Pty Ltd

AbbeyNeo™ Antibiotic Feed Additive

ACTIVE CONSTITUENT: 600g/kg NEOMYCIN (as Sulfate)



Broad-spectrum aminoglycoside antibiotic for the treatment of bacterial scours caused by organisms sensitive to Neomycin, including Neomycin sensitive strains of *E. coli*, *Salmonella* spp., *Pseudomonas* spp., in poultry, pigs, and cattle.

DIRECTIONS FOR USE

DOSAGE AND ADMINISTRATION

BLEND WITH A SMALL QUANTITY OF FEED BEFORE MIXING INTO LARGE QUANTITIES.

The recommended dose rate is 8-22 mg Neomycin / kg bodyweight / day. The inclusion rate of Neomycin in feed should be determined by the prescribing veterinarian. The minimum daily dose is 8mg Neomycin / kg bodyweight.

Use medicated feed for 3 to 5 days. If no improvement is seen after 3 days, the diagnosis should be re-established.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

MEAT: REMOVE ALL MEDICATED FEED for Broilers – 5 days; turkeys – 14 days; laying hens – 14 days; pigs – 20 days; cattle – 30 days before slaughter for human consumption.

MILK: Zero (0) days

EGGS: Zero (0) days

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

TRADE ADVICE

An ESI has not been established for this product. Note—observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Abbey Animal Health Pty Ltd on 02 8088 0720 before using this product or APVMA website (apvma.gov.au/residues).

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre: Phone Australia 131 126

DISPOSAL

Single-rinse or Shake and empty containers into medicated feed. Do not dispose of undiluted chemicals on site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or products.

STORAGE

Store below 30°C (room temperature). Store in a dry place.

APVMA Approval Number: 67805/140221