









# EQUIMATE™ ORAL PROGESTAGEN FOR HORSES

ACTIVE CONSTITUENT: 2.2mg/mL ALTRENOGEST

## Your teammate when it comes to breeding

For regulation and control of the breeding cycle of mares and the maintenance of pregnancy in habitually aborting mares, or mares at risk of early abortion.

-  Suppress oestrus
-  Induce oestrus
-  Maintain pregnancy
-  Use strategically to accommodate breeding or performance schedule
-  Easy to administer



	EQUIMATE™	Regu-mate®
Active Constituent	2.2mg/mL ALTRENOGEST	2.2mg/mL ALTRENOGEST
Species	Horses	Horses
Route of Administration	Oral	Oral
<b>Dose Rate</b>		
Induction of oestrus	1mL/50kg for 10 consecutive days	1mL/50kg for 10 consecutive days
Suppression of oestrus	1mL/50kg for 15 consecutive days	1mL/50kg for 15 consecutive days
Control Breeding Cycle	1mL/50kg for 15 consecutive days	1mL/50kg for 15 consecutive days
Maintenance of pregnancy	1mL/50kg from 2 <sup>nd</sup> or 3 <sup>rd</sup> day post ovulation for 120 days	1mL/50kg from 2 <sup>nd</sup> or 3 <sup>rd</sup> day post ovulation for 120 days

EQUIMATE™ is a trademark of Abbey Laboratories Pty Ltd

Regu-Mate® is a registered trademark of Intervet International B.V

# EQUIMATE™ ORAL PROGESTAGEN FOR HORSES

ACTIVE CONSTITUENT: 2.2mg/mL ALTRENOGEST



## For regulation and control of the breeding cycle of mares and the maintenance of pregnancy in habitually aborting mares, or mares at risk of early abortion.

### INDICATIONS

For regulation and control of the breeding cycle of mares. To induce ovulatory oestrus early in the breeding season in mares where some follicular activity exists. For the suppression of oestrus either during prolonged oestrus or in normally cycling mares. For the control of the ovarian cycle in breeding mares to allow the most efficient use of the stallion.

For the maintenance of pregnancy in habitually aborting mares, or mares at risk of early embryonic death or abortion, where progesterone deficiency is thought to be the underlying cause.

### DIRECTIONS FOR USE

#### Contraindications

DO NOT USE in male animals.

This product is contraindicated for use in mares suffering from uterine infection.

#### Precautions

Unused feed must be destroyed and not given to any other animals.

EQUIMATE™ must be added to the feed immediately prior to consumption and not stored.

### DOSAGE AND ADMINISTRATION

For oral administration.

Administer 1mL EQUIMATE™ per 50kg bodyweight (equivalent to 0.044mg Altrenogest per kg) daily.

#### 1. Regulation and control of the breeding cycle of mares:

- a. For the induction of ovulatory oestrus or to induce cyclical ovarian activity in mares with some follicle activity early in the breeding season: Administer EQUIMATE™ for 10 consecutive days.

- b. For suppression of oestrus in normally cycling mares: Administer EQUIMATE™ daily for 15 consecutive days.

For suppression of prolonged oestrus: Administer EQUIMATE™ for 10 consecutive days.

- c. To control the breeding cycle of mares to allow efficient use of stallion: Administer EQUIMATE™ daily for 15 consecutive days.

#### 2. For the treatment of habitually aborting mares, or mares at risk of early embryonic death or abortion:

Administer EQUIMATE™ daily, from the 2nd or 3rd day after ovulation in the mated or inseminated mare, continuing until day 120 of gestation.

#### Note:

Ultrasound scanning to confirm pregnancy is recommended on day 14 or 15 after ovulation. If the mare is non-pregnant it is advised to cease EQUIMATE™ administration and to administer an injection of Prostaglandin F2 $\alpha$  to destroy any luteal tissue that may be still present, allowing the mare to return to oestrus and be re-mated.

### MEAT WITHHOLDING PERIODS (HORSES)

**DO NOT USE less than 28 days before slaughter for human consumption.**

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

### ADDITIONAL USER SAFETY INFORMATION

Care should be taken to avoid contact between the solution and women of the childbearing age.

Additional information is listed in the safety data sheet (SDS).

### DISPOSAL

Dispose of container by wrapping with paper and putting in garbage.

### STORAGE

Store below 25 °C (air conditioning). Protect from light.

**APVMA Approval Number:  
88398/122367**

EQUIMATE™ is a trademark of Abbey Laboratories Pty Ltd