

# EXITIX™ POUR ON TICK DEVELOPMENT INHIBITOR

ACTIVE CONSTITUENT: 25g/L FLUAZURON

Solvents: 100g/L N-(N-OCTYL)-2-PYRROLIDONE, 100g/L N-(N-DODECYL)-2-PYRROLIDONE, 415g/L N-METHYL-2-PYRROLIDONE

## Tick protection made easy

For strategic control of Fluaazuron susceptible cattle tick *Rhipicephalus microplus*, including strains resistant to organophosphates, synthetic pyrethroids and amidines, on beef cattle.

- 🌿 Tick development inhibitor
- 🌿 Breaks the tick life cycle
- 🌿 Ready to use, pour on formulation
- 🌿 Cost Effective
- 🌿 Pack sizes: 5L Backpack, 20L



	EXITIX™ POUR ON	Acatak®
Active Constituent	25g/L FLUAZURON	25g/L FLUAZURON
Species	Cattle	Cattle
Dose Rate	3mL/50kg bodyweight	3mL/50kg bodyweight
Route of Administration	Pour on	Pour on
Withholding periods		
Meat	42 days	42 days
Milk	DO NOT USE	DO NOT USE
ESI	42 Days	42 Days

EXITIX™ is a trademark of Abbey Laboratories Pty Ltd

Acatak® is a registered trademark of Elanco Tiergesundheits AG

# EXITIX™ POUR ON TICK DEVELOPMENT INHIBITOR

ACTIVE CONSTITUENT: 25g/L FLUAZURON

Solvents: 100g/L N-(N-OCTYL)-2-PYRROLIDONE, 100g/L N-(N-DODECYL)-2-PYRROLIDONE, 415g/L N-METHYL-2-PYRROLIDONE



## For strategic control of Fluazuron susceptible cattle tick *Rhipicephalus microplus*, including strains resistant to organophosphates, synthetic pyrethroids and amidines on beef cattle.

This product contains Fluazuron. It is effective against Fluazuron sensitive cattle ticks on beef cattle. Resistance may develop to any chemical.

### DIRECTIONS FOR USE

#### Restraint

DO NOT USE on cattle which are producing or may in future produce milk that may be used or processed for human consumption.

#### Contraindication

This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.

### DOSAGE AND ADMINISTRATION

Animals and Pests	States	Bodyweight (kg)	Dose Volume	Cattle treated by 5L	Cattle treated by 20L
Beef cattle after weaning: Cattle tick	QLD, NT, WA and NSW only	101-150	9	555	2222
		151-200	12	416	1666
		201-250	15	333	1333
		251-300	18	277	1111
		301-350	21	238	952
		351-400	24	208	833
		401-450	27	185	740
		451-500	30	166	666
		501-550	33	151	606
		551-600	36	138	555
		601-650	39	128	512
		Over 650		3mL/50kg bodyweight	

### CRITICAL COMMENTS

- Retreat when you first see ticks but not before at least last 6 weeks after the previous treatment.
- Maximum number of treatments per annual tick season: 3.
- Apply in 2 bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with applicator.
- The weight of cattle should be determined by either scales or weighband. Dose individuals according to the heaviest animal in the mob. DO NOT underdose.
- There is no need to treat calves for cattle tick as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.
- This product should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.

7. This product is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of this product commences within three days of treatment.

8. Not suitable for cleaning cattle moving into "gazetted" cattle tick free areas.

### GENERAL DIRECTIONS

**Product treated areas - introduction of new cattle:** Tick infested cattle introduced onto properties where this product treatment regime is in place should be treated with EXITIX™ prior to release.

**Handling of treated animals:** Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.

For local tick control recommendations contact your local department of agriculture.

### Cleaning Instructions

After use wash applicator with detergent and hot water. Dispose of the rinsate in a disposal pit away from desirable plants and their roots and watercourses.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 42 days before slaughter for human consumption.**

**Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.**

**MILK: DO NOT USE on cattle which are producing or may in the future produce milk that may be used or processed for human consumption.**

### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal health Pty Ltd on 020 8088 0720 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**

### SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes and skin. Will irritate the eyes. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders.

When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately.

If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking and smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. Additional information is listed in Safety Data Sheet.

### ENVIRONMENTAL PROTECTION

This product is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the product or used containers.

### DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### STORAGE

Store below 30°C (Room Temperature). Store in original container tightly closed in a dry, cool place. Store out of direct sunlight.

APVMA Approval Number: 90312/131146