

# EXITIX™ GOLD

**BROAD SPECTRUM POUR ON FOR CATTLE**

ACTIVE CONSTITUENTS: 5g/L IVERMECTIN  
15g/L FLUAZURON



## Gold Standard in Tick Protection

For treatment and control of Ivermectin sensitive internal and external parasites of beef cattle. For strategic control of Flouazuron susceptible tick *Rhipicephalus microplus*, including strains resistant to organophosphates, synthetic pyrethroids and amidines, on beef cattle.




- **Broad spectrum parasiticide**
- **Tick development inhibitor**
- **Reduce the build up of tick population**
- **Persistent activity against buffalo fly**

EXITIX™ GOLD		ACATAK® DUOSTAR
ACTIVE CONSTITUENTS	5g/L IVERMECTIN, 15g/L FLUAZURON	5g/L IVERMECTIN, 15g/L FLUAZURON
SPECIES	Cattle	Cattle
DOSE RATE	5mL/50kg	5mL/50kg
ROUTE OF ADMINISTRATION	Pour On	Pour on
WITHHOLDING PERIODS		
MEAT: Cattle	28 days	28 days
MILK: Cattle	DO NOT USE	DO NOT USE
ESI	42 days	42 Days

EXITIX™ is a trademark of Abbey Laboratories Pty Ltd

ACATAK® is a registered trademark of Elanco Tiergesundheits AG



# EXITIX™ GOLD

## BROAD SPECTRUM POUR ON FOR CATTLE

ACTIVE CONSTITUENTS: 5g/L IVERMECTIN  
15g/L FLUAZURON

For treatment and control of Ivermectin sensitive internal and external parasites of beef cattle.

### INDICATIONS

This product contains Fluazuron and Ivermectin (a member of the avermectin (macrocyclic lactone) family). It is effective against cattle ticks and Ivermectin-sensitive internal and external parasites of beef cattle (listed below):

#### Cattle tick

*Rhipicephalus microplus*

#### Gastrointestinal roundworms

#### For the control of adult and immature (L<sub>4</sub>)

Small Brown Stomach Worm (including inhibited stages) *Ostertagia ostertagi*

Barber's Pole Worm *Haemonchus placei*

Stomach Hair Worm *Trichostrongylus axei*

Black Scour Worm *Trichostrongylus colubriformis*

Small Intestinal Worm *Cooperia* spp.

Nodule Worm *Oesophagostomum radiatum*,

*Oesophagostomum venulosum*

#### And adult stages of

Whipworm *Trichuris* spp.

Intestinal Threadworm *Strongyloides papillosus*

#### And immature stages of

Thin Necked Intestinal Worm L<sub>4</sub> *Nematodirus* spp.

#### Lungworm (adult and immature)

*Dictyocaulus viviparus*

#### Eyeworm (adult stages) *Thelazia* spp.

#### Sucking and Biting Lice *Linognathus vituli*

*Haematopinus eurysternus*

*Bovicola (Damalinia) bovis*

*Solenopotes capillatus*

#### Mites

*Chorioptes bovis*

*Sarcoptes scabiei* var. *bovis*

#### Buffalo flies

*Haematobia irritans exigua*

Resistance may develop to any chemical.

#### Persistent activity

Product applied at the recommended dose of 5mL/50kg liveweight controls infestations of *Haematobia irritans exigua* (buffalo fly) for up to 7 days after treatment.

### DIRECTIONS FOR USE

#### Restraint

DO NOT USE in cattle which are producing or may in future produce milk for human consumption.

#### Contraindication

This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.

### DOSAGE AND ADMINISTRATION

#### DOSE RATE TABLE:

Animals and pests	Bodyweight (kg)	Dose Volume (mL)
Cattle Tick	101-150	15
( <i>Rhipicephalus microplus</i> )	151-200	20
Ivermectin sensitive gastrointestinal roundworms	201-250	25
(including inhibited <i>Ostertagia ostertagi</i> ), lungworms, sucking and biting lice, sarcoptic and chorioptic mites, and buffalo fly	251-300	30
( <i>Haematobia irritans exigua</i> )	301-350	35
	351-400	40
	401-450	45
	451-500	50
	501-550	55
	551-600	60
	601-650	65
	over 650	5mL/50kg

#### CRITICAL COMMENTS

1. Retreat when you first see ticks but not before at least last 6 weeks after the previous treatment.
2. Maximum number of treatments per annual tick season: 3.
3. To achieve optimal efficacy, this product relies on a combination of absorption through skin and oral ingestion (via grooming). For oral ingestion it is critical that the product is applied as 2 x 7 cm wide bands, from shoulder to the rump, that are well separated on the other side of the spine. The EXITIX™ POUR ON applicator is recommended for use with this product.
4. The weight of cattle should be determined by either scales or weighband. Dose individuals according to the heaviest animal in the mob. Recommended dose rate gives 1.5mg Fluazuron and 0.5mg Ivermectin per kg bodyweight of beef cattle. DO NOT underdose.
5. There is no need to treat calves for cattle tick as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.
6. This product should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.
7. The effect of this product is not adversely affected if rain occurs after application.

#### GENERAL DIRECTIONS

##### Product treated areas- introduction of new cattle:

Tick infested cattle introduced onto properties where this product treatment regime is in place should be treated with EXITIX™ GOLD prior to release.

**Handling of treated animals:** Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.

To avoid operator contact with this product when concurrent treatments are carried out, this product should be applied last. For local tick control recommendations contact your local department of agriculture.

#### Cleaning Instructions

After use wash applicator with detergent and hot water. Dispose of the rinsate in a disposal pit away from desirable plants and their roots and watercourses.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 28 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.**

**MILK: DO NOT USE on cattle which are producing or may in the future produce milk that may be used or processed for human consumption.**

#### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal health Pty Ltd on 02 8088 0720 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**

#### SAFETY DIRECTIONS

Harmful if swallowed. May irritate the skin. Will irritate the eyes. Avoid contact with eyes and skin. When preparing product for use and using product, wear elbow-length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves.

#### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Additional information is listed in Safety Data Sheet.

#### ENVIRONMENTAL PROTECTION

This product is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the product or used containers.

#### DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### STORAGE

Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. Store out of direct sunlight.

EXITIX™ is a trademark of Abbey Laboratories Pty Ltd