










# FASITAK™ POUR ON FOR CATTLE

ACTIVE CONSTITUENTS: 300 g/L TRICLABENDAZOLE, 5 g/L ABAMECTIN  
Solvent: 500 G/L DIETHYLENE GLYCOL MONOBUTYL ETHER

## Give fluke the flick

For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.

-  Proven efficacy against early immature, immature and mature stages of liver fluke
-  Treats and controls a wide range of adult and immature stages of worms
-  Aids in control of lice, Buffalo Fly, Cattle Tick
-  Easy to use pour-on application; no hassle
-  Dual Actives
-  Cost Effective
-  Pack sizes: 2.5L Backpack, 10L



	FASITAK™ Pour On	FASIMEC® Cattle Pour On
Active Constituents	300 g/L TRICLABENDAZOLE 5 g/L ABAMECTIN	300 g/L TRICLABENDAZOLE 5 g/L ABAMECTIN
Species	Cattle	Cattle
Dose Rate	1mL/10kg	1mL/10kg
Route of Administration	Pour on	Pour on
<b>Withholding periods</b>		
Meat	49 days	49 days
Milk	DO NOT USE	DO NOT USE
ESI	140 Days	140 Days

FASITAK™ is a trademark of Abbey Laboratories Pty Ltd

FASIMEC® is a registered trademark of Elanco Tiergesundheit AG

# FASITAK™ POUR ON FOR CATTLE

ACTIVE CONSTITUENTS: 300 g/L TRICLABENDAZOLE, 5 g/L ABAMECTIN  
Solvent: 500 g/L DIETHYLENE GLYCOL MONOBUTYL ETHER



## For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.

FASITAK™ Pour-On contains abamectin, a member of the macrocyclic lactone family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains Triclabendazole, a member of the benzimidazole family of chemicals, which is effective against liver fluke. Resistance may develop to any chemical.

### INDICATIONS

**For the treatment and control of adult and immature stages of:**

#### Gastro-intestinal roundworms

Barber's Pole Worm *Haemonchus placei*

Small Brown Stomach Worm *Ostertagia*

*ostertagi* (including inhibited stages)

Stomach Hair Worm *Trichostrongylus axei*

Small Intestinal Worm *Cooperia oncophora*, *C. punctata*

Hookworm *Bunostomum phlebotomum*

Thin-necked Intestinal Worm *Nematodirus*

*spathiger*

Intestinal Threadworm *Strongyloides papillosus*

Large Bowel Worm *Oesophagostomum radiatum*

Whipworm *Trichuris* spp.

**Lungworm** *Dictyocaulus viviparus*

**Liver Fluke** *Fasciola hepatica* (including adult (12 weeks old), immature (6 weeks old) and early immature (4 weeks old) larvae)

#### Sucking and Biting Lice

Long nosed sucking Lice *Linognathus vituli*

Short nosed sucking Lice *Haematopinus*

*eurytjernus*

Biting Lice *Bovicola bovis*

Little sucking Lice *Solenopotes capillatus*

#### Aids in control of:

**Buffalo Fly** *Haematobia irritans exigua*

**Cattle Tick** *Boophilus microplus* (Not recommended in strategic dipping programs for cattle tick control)

Resistance may develop to any chemical.

### DIRECTIONS FOR USE

#### Restraint

DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption.

#### Contraindication

This product is contraindicated for use in calves under 50kg bodyweight.

#### DOSAGE AND ADMINISTRATION

FASITAK™ POUR-ON dose rate is 1mL/10kg bodyweight. Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

Liveweight (kg)	Dose Volume (mL)	10L Treats	2.5L Treats
51-75	7.5	1330	333
76-100	10	1000	250
101-125	12.5	800	200
126-150	15	600	166
151-175	17.5	570	142
176-200	20	500	125
201-250	25	400	100
251-300	30	330	83
301-350	35	280	71
351-400	40	250	62
401-450	45	220	55
451-500	50	200	50
501-550	55	180	45
551-600	60	160	41
601-650	65	150	38
Over 650kg dose at rate of 1mL/10kg bodyweight.			

A representative sample of animals should be weighed either with scales or a weighband before treatment. Dose according to the heaviest animal by live weight in the group (cows, bulls, heifers, etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight to avoid excessive overdosing. Check the accuracy of guns regularly.

FASITAK™ POUR-ON is rainfast. Rain occurring as early as 2 hours after treatment is unlikely to affect the efficacy of the product.

A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. Do not brush off or wash off. This deposit may last for a few days or weeks depending on weather conditions.

Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian.

After use clean all equipment with cooking oil. Do not use water.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 49 days before slaughter for human consumption. Beef calves born to cows that were treated with FASITAK™ POUR ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the dam.**

**MILK: DO NOT USE in lactating cows which are producing or may in future produce milk that may be used or processed for human consumption.**

#### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 140 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)). Beef calves born to cows that were treated with FASITAK™ POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for export for 140 days after treatment of the dam.**

#### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, first quickly, gently, and completely blot or wipe chemical from eyes then flush with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing, quickly and gently blot or wipe away chemical with a tissue or cloth, preferably impregnated with cooking oil, then wash skin thoroughly using soap and water. Do not use water first. Additional information is listed in Safety Data Sheet.

#### ENVIRONMENTAL PROTECTION

Very toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

#### DISPOSAL

Triple or (preferably) pressure rinse the container and dispose of rinsate in compliance with relevant local, state or territory government. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### STORAGE

Store below 30°C (Room Temperature).

APVMA Approval Number: 87638/128618

FASITAK™ is a trademark of Abbey Laboratories Pty Ltd