

LINCO-SPEC™ PREMIX



**ACTIVE CONSTITUENTS: 22 g/kg SPECTINOMYCIN (as Spectinomycin sulphate),
22 g/kg LINCOMYCIN (as Lincomycin hydrochloride)**

Dual active efficacy

For treatment and control of swine dysentery (Vibronic dysentery, bloody scours); salmonellosis (necrotic enteritis); enteritis associated with *E.coli* and as an aid in control of mycoplasmal pneumonia.

- **Increased efficacy with the combination of 2 actives**
- **Controls key bacteria in swine**
- **Easy to use formulation**
- **24-hour WHP**



	LINCO-SPEC™ PREMIX	LINCO-SPECTIN® PREMIX
ACTIVE CONSTITUENT	22 g/kg SPECTINOMYCIN (as Spectinomycin sulphate), 22 g/kg LINCOMYCIN (as Lincomycin hydrochloride)	22 g/kg SPECTINOMYCIN (as Spectinomycin sulphate), 22 g/kg LINCOMYCIN (as Lincomycin hydrochloride)
SPECIES	Pigs	Pigs
DOSE RATE	1kg LINCO-SPEC™ per tonne of complete feed to give 44g of antibiotic activity	1kg LINCO-SPECTIN® per tonne of complete feed to give 44g of antibiotic activity
WITHOLDING PERIODS		
Meat	1 day	1 day

LINCO-SPEC™ is a trademark of Abbey Laboratories Pty Ltd.

LINCO-SPECTIN® is a registered trademark of Zoetis Australia Pty Ltd



LINCO-SPEC™ PREMIX

ACTIVE CONSTITUENTS:

22 g/kg SPECTINOMYCIN (as Spectinomycin sulphate)

22 g/kg LINCOMYCIN (as Lincomycin hydrochloride)

For treatment and control of swine dysentery (Vibronic dysentery, bloody scours); salmonellosis (necrotic enteritis); enteritis associated with *E.coli* and as an aid in control of mycoplasmal pneumonia.

DIRECTIONS FOR USE

Restraint

DO NOT continue medication beyond the period prescribed by the veterinarian.

Precautions

For use in pigs only. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feed containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Lincomycin should not be used in animals known to be hypersensitive to the product. Do not use the water treatment and feed treatment simultaneously.

Side Effects

Excessive dosage may lead to an increased incidence of side effects which may include rectal prolapse.

Occasionally, swine treated with Lincomycin, within the first two days after the onset of treatment, may develop diarrhoea and/or swelling of the anus. On rare occasions, some pigs may show reddening of skin and irritable behaviour. These conditions have been self-correcting within 5 to 8 days without discontinuing Lincomycin treatment.

DOSAGE AND ADMINISTRATION

For the treatment and control of diseases specified, complete feeds containing 44 grams of antibiotic activity per tonne should be fed.

GENERAL DIRECTIONS

Mixing Directions

LINCO-SPEC™ PREMIX must be thoroughly mixed in feeds before use.

For making medicated complete feeds, add 1 kg LINCO-SPEC™ PREMIX per tonne of complete feed to give 44 g of antibiotic activity per tonne.

Medicated feed should constitute the sole ration of treated and growing swine.

Good manufacturing practices should be observed in preparing feeds containing LINCO-SPEC™ PREMIX. This includes appropriate clean-out procedures to avoid cross contamination.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 1 day before slaughter for human consumption.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact Abbey Animal Health Pty Ltd on 02 8088 0720.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

Additional information is listed in the Safety Data Sheet.

DISPOSAL

Shake container into medicated feed, water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. Break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (Room Temperature).

PROTECT FROM MOISTURE. Store opened bag in a dry place to prevent caking.

May cause an allergic skin reaction.

May cause damage to organs through prolonged or repeated exposure.

May cause allergy or asthma symptoms or breathing difficulties if inhaled.

In case of inadequate ventilation wear respiratory protection.

Do not breathe dust.

If experiencing respiratory symptoms: Call a POISON CENTRE or doctor/physician

IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing.

LINCO-SPEC™ is a trademark of Abbey Laboratories Pty Ltd

APVMA Approval No. 86922/117062