



# SULFA-B™ Oral Paste Antibiotic Mucolytic for Horses

ACTIVE CONSTITUENTS: 325mg/mL SULFADIAZINE, 65mg/mL TRIMETHOPRIM,  
20mg/mL BROMHEXINE HYDROCHLORIDE

New  
Pack  
Size

## Peace of mind when you can't be there for every dose

For the treatment of productive upper respiratory tract infections in horses.

- Antibiotic and mucolytic action
- Reduced recovery time for acute respiratory infections
- Significant blood levels achieved within 30 minutes
- Ready to use dial-a-dose syringes



|                         | SULFA-B™ Oral Paste   | Bromotrimidine Paste  |
|-------------------------|---|---|
| Active Constituents     | 325mg/mL Sulfadiazine,<br>65mg/mL Trimethoprim,<br>20mg/mL Bromhexine HCl | 450mg/g Sulfadiazine,<br>90mg/g Trimethoprim,<br>9mg/g Bromhexine HCl |
| Species                 | Equine  | Equine  |
| Dose rate               | 5mL/100kg twice a day,<br>12 hourly for 5 days                            | 20mL/450kg twice daily  |
| Route of Administration | Oral  | Oral  |

SULFA-B™ is a trademark of Abbey Laboratories Pty Ltd

International Animal Health Products Pty Ltd

# SULFA-B™ Oral Paste Antibiotic Mucolytic for Horses

ACTIVE CONSTITUENTS: 325mg/mL SULFADIAZINE, 65mg/mL TRIMETHOPRIM,  
20mg/mL BROMHEXINE HYDROCHLORID



## For the treatment of productive upper respiratory tract infections in horses.

### INDICATIONS

SULFA-B™ is indicated for the treatment of productive upper respiratory tract infections in horses where those infections are susceptible to Sulfadiazine potentiated with Trimethoprim and where the mucolytic effect of Bromhexine HCl is desired.

The Sulfonamide acts by blocking the biosynthesis of folic acid in the bacterial cell and the Trimethoprim interferes in the reduction process of folic acid in the bacterial cell to the biologically active form tetrahydrofolate. Bromhexine HCl aids in the removal of the thick, tenacious mucoid secretions in the airways by reducing viscosity of the acid mucopolysaccharide fibers from the mucous glands in nose, throat and lungs. This facilitates the availability of the antibacterial agents to the infected areas to reduce recovery time following acute respiratory infection.

### DIRECTIONS FOR USE

#### Contraindications

SULFA-B™ is generally well tolerated with a low incidence of adverse reactions. Do not use in renal or hepatic dysfunction or with known previous hypersensitivity to Sulfonamides. Take care with gut superinfection which can lead to toxic megacolon. The animal should have access to food and cool, clean water. Discontinue treatment if there is marked inappetence or signs of abdominal discomfort. Long term therapy may require replenishing treatment with multi-vitamins, vitamin B12 and folic acid supplements.

### DOSAGE AND ADMINISTRATION

Horses: 5mL/100kg twice a day 12 hourly for 5 days. A 30mL syringe will deliver the dose once a day for horse weighing 600kg at the dose rate of 5mL/100kg.

#### For 30mL syringe:

| Weight of Horse (kg) | Number of syringes required per day |
|----------------------|-------------------------------------|
| 300                  | 1                                   |
| 600                  | 2                                   |

To administer, remove cap from the syringe and direct the nozzle of the syringe into the side of the horse's mouth. Depress the plunger to dispense the desired dose towards the back of the tongue. Keep the mouth closed for a few seconds to make sure the dose is swallowed. Replace cap after use.

### GENERAL DIRECTIONS

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

If used in performance animals, the regulations of the relevant authorities regarding medication should be observed.

### WITHHOLDING PERIODS

**MEAT WITHHOLDING PERIODS (HORSES): DO NOT USE less than 28 days before slaughter for human consumption.**

**Any variation by the prescribing veterinarian to the approved use pattern, may require an extended withholding period.**

### EXPORT SLAUGHTER INTERVAL (ESI)

**An ESI has not been established for this product.**

**Note – observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Abbey Animal Health Pty Ltd on 02 8088 0720 before using this product.**

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### ADDITIONAL USER SAFETY

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands thoroughly after use. Flush mouth liberally if accidental ingestion.

Additional information is listed in the safety data sheet (SDS).

### ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemical or used containers.

### DISPOSAL

Dispose of container by wrapping with paper and putting in garbage.

### STORAGE

Store below 25° C (air conditioning). Protect from light.

**APVMA Approval Number:  
49953/138984**