

# SELECTAN®

SOLUTION FOR INJECTION FOR CATTLE AND PIGS

ACTIVE CONSTITUENT: 300mg/mL FLORFENICOL



## SELECT the best

For the treatment of bacterial infections in cattle and pigs susceptible to florfenicol.


- **Single product for cattle and swine**
- **Low viscosity = Excellent syringeability**
- **Easy to inject, even at low temperatures**
- **Quicker to inject allowing more animals to be treated in less time**
- **Plastic bottle to prevent breakages**
- **Shorter Meat WHP for cattle**



	SELECTAN®	NUFLOR® LA
<b>ACTIVE CONSTITUENT</b>	300mg/mL Florfenicol	300mg/mL Florfenicol
<b>SPECIES</b>	Cattle and Pigs	Cattle and Pigs
<b>I. M. DOSE RATE</b>		
Cattle	1mL/15kg	1mL/15kg
Pigs	0.75mL/15kg	0.75mL/15kg
<b>WITHHOLDING PERIODS</b>		
Meat: Cattle	28 days	42 days
Milk: Cattle	Do not use	Do not use
Meat: Pigs	12 days	12 days
<b>ESI</b>		
Cattle	42 days	42 days
Pigs	18 days	12 days

SELECTAN® is a registered trademark of Laboratoris Hipra, S.A.

NUFLOR® is a registered trademark of Intervet international B.V.



# SELECTAN<sup>®</sup>

SOLUTION FOR INJECTION  
FOR CATTLE AND PIGS

## ACTIVE CONTITUENT: 300mg/mL FLORFENICOL

For the treatment of bacterial infections in cattle and pigs susceptible to florfenicol.

### DIRECTIONS FOR USE

#### Restraints

Cattle and Pigs: DO NOT repeat the treatment schedule of two (2) doses per animal, administered 48 hours apart.

Cattle: DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE in calves to be processed for veal.

DO NOT exceed 10mL at any one injection site (intramuscular).

Pigs: DO NOT exceed 2mL at any one injection site (intramuscular).

#### Contraindications

This product should not be used in cattle or pigs intended for breeding since the effect of florfenicol on bovine and porcine reproductive performance and pregnancy has not been assessed.

#### Side Effects

Mild, transient inappetance, decreased water consumption or diarrhoea may occur in cattle following treatment. Transient diarrhoea and/or perianal erythema/swelling may occur in pigs following treatment. These symptoms should return to normal shortly after treatment. As seen with many antibiotics high dosage and/or prolonged administration may lead to superinfections by non-sensitive organisms such as yeasts and fungi.

In cattle injection may result in local tissue reaction which persists beyond 35 days and may result in trim loss of edible tissue at slaughter.

In pigs injection may result in local tissue reaction which persists for up to 21 days and may result in trim loss of edible tissue at slaughter.

Tissue reactions at injection sites other than the neck are likely to be more severe.

### DOSAGE AND ADMINISTRATION

Use the contents within 28 days of first broaching of the bottle. Discard unused portion.

**Cattle:** Intramuscular - 20mg/kg of bodyweight (1mL per 15kg). A second dose should be given 48 hours later. Do not exceed 10mL at any one injection site. Inject into muscle tissue on the side of the neck.

**Pigs:** Intramuscular - 15mg/kg bodyweight (0.75mL per 15kg). Two injections are given 48 hours apart. Do not exceed 2mL at any one injection site. Inject into muscle tissue on the side of the neck.

MANUFACTURED BY: LABORATORIOS HIPRA, S.A.

DISTRIBUTED BY:

**ABBEY ANIMAL HEALTH PTY LTD**

16 VOYAGER CIRCUIT

GLENDENNING NSW 2761

INFO LINE 02 8088 0720

[www.abbeylabs.com.au](http://www.abbeylabs.com.au)

### GENERAL DIRECTIONS

Florfenicol is a synthetic, broad spectrum antibiotic active against many gram-negative and gram-positive bacteria isolated from domestic animals. Florfenicol acts by inhibiting bacterial protein synthesis at the ribosomal level.

Clinical improvement should be evident in most treated subjects within 24 hours of initiation of therapy. If a positive response is not seen within 24 hours after the second injection, the diagnosis should be reconfirmed.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

### WITHHOLDING PERIODS

**Cattle: MEAT: 28 days. MILK: DO NOT USE in cows which are/may produce milk used or processed for human consumption.**

**Pigs: MEAT: 12 days**

### TRADE ADVICE - EXPORT SLAUGHTER INTERVALS (ESI)

**Cattle: 42 days.**

**Pigs: 18 days**

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131 126).

### DISPOSAL


Dispose of container by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labeled 'sharps' container.

### STORAGE

Store below 25°C (Air conditioning). Protect from light. Do not freeze.

**SELECTAN<sup>®</sup> is a registered trademark of Laboratorios Hipra, S.A.**

APVMA Approval No: 68700/103531



**SELECTAN<sup>®</sup>**