







# TYLOVET™ TYLOSIN TARTRATE SOLUBLE

ACTIVE CONSTITUENT: 800mg/g TYLOSIN (as the tartrate)

## A watertight defence against infection

TYLOVET™ Tylosin Tartrate Soluble is a water-soluble macrolide antibiotic product.

As an aid in the prevention and treatment of Chronic Respiratory Disease in broilers and replacement chickens, infectious sinusitis in turkeys and swine dysentery in pigs.

-  Easy to use - just mix into water!
-  Preventative or therapeutic dose options
-  Treats bacterial and respiratory infection
-  High safety margin



TYLOVET™ SOLUBLE	
Active Constituent	800mg/g TYLOSIN as the tartrate
Species	Broilers, Chickens, Turkeys, Pigs
Route of Administration	Water soluble
Dose rate	100g/5L and further diluted as necessary
Pigs	Diluted to 400L
Chickens and Turkeys	Diluted to 200L
Withholding periods	
Meat	Nil
Eggs	DO NOT USE

TYLOVET™ is a trademark of Abbey Laboratories Pty Ltd

# TYLOVET™ TYLOSIN TARTRATE SOLUBLE

ACTIVE CONSTITUENT: 800mg/g TYLOSIN (as the tartrate)



**TYLOVET™ Tylosin Tartrate Soluble is a water-soluble macrolide antibiotic product.**

**As an aid in the prevention and treatment of Chronic Respiratory Disease in broilers and replacement chickens, infectious sinusitis in turkeys and swine dysentery in pigs.**

## **DIRECTIONS FOR USE**

### **Restraint**

DO NOT USE in birds, which are producing or may in the future produce eggs or egg products for human consumption.

### **Mixing Directions**

Add at the rate of 100g to 5 litres water and dissolve thoroughly. Further dilute this concentrated solution as per the dosage recommended below. Prepare a fresh TYLOVET™ solution every three days.

## **DOSAGE AND ADMINISTRATION**

**PIGS:** As an aid in the treatment of swine dysentery. Further dilute the concentrated solution to provide 400 litres of medicated drinking water and treat for 3 to 10 days. Treatment should be continued until 24 hours after the cessation of symptoms.

**CHICKENS AND TURKEYS:** Further dilute the concentrated solution to provide 200 litres of medicated drinking water. Provide as the only source of drinking water.

**CHICKENS:** As an aid in the prevention and treatment of chronic respiratory disease in broilers and replacement chickens.

**PREVENTATIVE:** Treat for the first three days of life. Repeat treatment for one day at 3 to 4 weeks of age, or to coincide with live virus vaccine or other stress.

**THERAPEUTIC:** Treat for 3-6 days depending on the severity of infection.

**TURKEYS:** As an aid in the treatment of infectious sinusitis. Treat for 2 to 5 days depending on the severity of infection.

## **WITHHOLDING PERIODS**

**MEAT: Zero (0) days.**

**Any variation by the prescribing veterinarian to the approved use pattern, may require an extended withholding period.**

**EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product.**

**Note – observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Abbey Animal Health Pty Ltd on 02 8088 0720 before using this product.**

## **SAFETY DIRECTIONS**

Repeated exposure may cause allergic disorders. Avoid contact with skin. Do not inhale dust when opening the container and preparing product for use and mixing. Wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and disposable dust face mask covering mouth and nose. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

## **DISPOSAL**

Shake container into medicated water. Do not dispose of undiluted chemicals on-site. Break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## **STORAGE**

Store below 25°C (air conditioning) in a dry place.

**APVMA Approval Number:  
54322/141865**

TYLOVET™ is a trademark of Abbey Laboratories Pty Ltd