








# TETRAMAX™ 100 INJECTION

ACTIVE CONSTITUENT: 96mg/mL OXYTETRACYCLINE (as the hydrochloride)

## Short acting oxytet

Broad spectrum antibiotic injection for use in cattle, sheep and pigs.

-  Short acting oxytetracycline
-  Broad spectrum, effective against both gram positive and gram negative bacteria
-  Flexible administration, can be used IM or slow IV
-  Multi species registration; cattle, sheep and pigs



	TETRAMAX™ 100	ALAMYCIN® SA 100
Active Constituent	96mg/mL OXYTETRACYCLINE (as the Hydrochloride)	100mg/mL OXYTETRACYCLINE
Species	Cattle, Sheep & Pigs	Cattle, Sheep & Pigs
Dose Rate: Cattle	2mL/50kg daily for 3-5 days	2mL/50kg daily for 3 - 5 days
Dose Rate: Sheep & Pigs	2-4.5mL/50kg daily for 3-5 days	2-4.5mL/50kg daily for 3-5 days
Route of Administration	IM or Slow IV	IM or Slow IV
<b>Withholding Periods</b>		
Meat: Cattle	11 days	11 days
Milk: Cattle / Ewes	60 hours (5 milkings)	60 hours (5 milkings)
Meat: Pigs	8 days	8 days
Meat: Sheep	15 days	15 days
<b>Export Slaughter Intervals</b>		
Cattle	11 days	11 days
Pigs	8 days	8 days
Sheep	Not established	Not established

TETRAMAX™ is a trademark of Abbey Laboratories Pty Ltd.

Alamycin® is a registered trademark of Norbrook Laboratories Ltd.

# TETRAMAX™ 100 INJECTION

ACTIVE CONSTITUENT: 96mg/mL OXYTETRACYCLINE (as the hydrochloride)



## Broad spectrum antibiotic injection for use in cattle, sheep and pigs.

### DIRECTIONS FOR USE

#### DOSAGE AND ADMINISTRATION

**Cattle:** 4mg/kg (2mL per 50kg bodyweight) daily for 3-5 days.

**Sheep and Pigs:** 4-9 mg/kg (2-4.5mL per 50kg bodyweight) daily for 3-5 days.

Intramuscular or slow intravenous injection.

If volume to be given is greater than 20mL it should be divided and injected into two sites. When kept at recommended storage temperatures, the product remains stable for 14 days after first broaching of the container.

#### WITHHOLDING PERIODS

**MEAT (CATTLE): DO NOT USE less than 11 days before slaughter for human consumption.**

**MEAT (PIGS): DO NOT USE less than 8 days before slaughter for human consumption.**

**MEAT (SHEEP): DO NOT USE less than 15 days before slaughter for human consumption.**

**MILK:** Milk collected from cows or ewes within 60 hours (5 milkings) following treatment **MUST NOT BE USED** for human consumption or processing. This milk should not be fed to bobby calves.

**Any variation by prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may result in the need to extend the approved withholding period.**

#### TRADE ADVICE- EXPORT SLAUGHTER INTERVAL (ESI)

**CATTLE: DO NOT USE less than 11 days before slaughter for export.**

**PIGS: DO NOT USE less than 8 days before slaughter for export.**

**SHEEP: This product does not have an ESI established.**

**Before using this product, confirm the current ESI with Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website [www.apvma.gov.au/residues/](http://www.apvma.gov.au/residues/).**

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

Additional information is listed in Safety Data Sheet.

#### DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

#### STORAGE

Store below 25°C (Air conditioning). Protect from light. When kept at the recommended storage temperatures, the product remains stable for 14 days after first broaching of the container.

**TetraMax™ is a trademark of Abbey Laboratories Pty Ltd.**

**APVMA Approval Number: 88577/121731**