

TYLOVET™

TYLOSIN TARTRATE SOLUBLE

ACTIVE CONSTITUENT: NOT LESS THAN 800mg/g TYLOSIN AS THE TARTRATE



Your Animals will TyLOVEt it

As an aid in the prevention and treatment of chronic respiratory disease in broilers and replacement chickens, infectious sinusitis in turkeys and swine dysentery in pigs

- Easy to use - Just mix into feed!
- Treats bacterial infections
- High safety margin
- Treats respiratory infections
- Cost Effective



TYLOVET	
ACTIVE CONSTITUENT	Not less than 800mg/g TYLOSIN as the tartrate
SPECIES	Broilers, Chickens, Turkeys, Pigs
DOSE RATE	100 g/5L and further diluted as necessary
Pigs	Diluted to 400 litres
Chickens and Turkeys	Diluted to 200 litres
ROUTE OF ADMINISTRATION	Water soluble
WITHHOLDING PERIODS	
Meat	NIL
Eggs	DO NOT USE

TYLOVET™ is a trademark of Abbey Laboratories Pty Ltd



TYLOVET™

TYLOSIN TARTRATE SOLUBLE

ACTIVE CONSTITUENT: NOT LESS THAN 800 MG/G TYLOSIN AS THE TARTRATE

Water Soluble Antimicrobial

As an aid in the Prevention and Treatment of Chronic Respiratory Disease in Broilers and Replacement Chickens, Infectious Sinusitis in Turkeys and Swine Dysentery in Pigs.

DIRECTIONS FOR USE

Restraint: DO NOT USE in birds, which are producing or may in the future produce eggs or egg products for human consumption.

Mixing Directions

Add at the rate of 100g to 5 litres water and dissolve thoroughly. Further dilute this concentrated solution as per the dosage recommended below. Prepare a fresh TYLOVET™ solution every three days.

DOSAGE AND ADMINISTRATION

PIGS: As an aid in the treatment of swine dysentery. Further dilute the concentrated solution to provide 400 litres of medicated drinking water and treat for 3 to 10 days. Treatment should be continued until 24 hours after the cessation of symptoms.

CHICKENS AND TURKEYS: Further dilute the concentrated solution to provide 200 litres of medicated drinking water. Provide as the only source of drinking water.

CHICKENS: As an aid in the prevention and treatment of chronic respiratory disease in broilers and replacement chickens.

PREVENTATIVE: Treat for the first three days of life. Repeat treatment for one day at 3 to 4 weeks of age, or to coincide with live virus vaccine or other stress.

THERAPEUTIC: Treat for 3-6 days depending on the severity of infection.

TURKEYS: As an aid in the treatment of infectious sinusitis. Treat for 2 to 5 days depending on the severity of infection.

WITHHOLDING PERIODS

MEAT: NIL

EGGS: DO NOT USE in birds, which are producing or may in the future produce eggs or egg products for human consumption.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may result in the need to extend the approved withholding period.

DISPOSAL

Shake and empty contents into medicated water. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

STORAGE

Store below 25°C (Air Conditioning) in a dry place.

TYLOVET™ is a trademark of Abbey Laboratories Pty Ltd

APVMA Approval No.54322/0601