


VB12™ PLUS SELENIUM FOR SHEEP AND CATTLE

ACTIVE CONSTITUENTS:
2mg/mL Hydroxocobalamin
(as Hydroxocobalamin acetate Vitamin B12)
4mg/mL Selenium as sodium selenate

B12 + Selenium in one convenient injection

For the treatment and control of Vitamin B12 and selenium deficiency in sheep and cattle.

- Contains Hydroxocobalamin, a longer acting form of B12
- Subcutaneous injection
- Can be used in sheep and cattle
- Cost effective pillow pack
- NIL WHP or ESI



	VB12™ PLUS SELENIUM	COBALIFE® VB12 PLUS SELENIUM
ACTIVE CONSTITUENTS	2mg/mL Hydroxocobalamin 4mg/mL Selenium	2mg/mL Hydroxocobalamin 4mg/mL Selenium
SPECIES	Sheep and Cattle	Sheep and Cattle
DOSE RATE		
Lambs	0.5mL at marking or weaning	0.5mL at marking or weaning
Sheep	1-1.5mL pre lambing	1-1.5mL pre lambing
Calves	2-3mL from 2 months of age	2-3mL from 2 months of age
Cows	4-6mL pre calving	4-6mL pre calving
ROUTE OF ADMINISTRATION	Subcutaneous Injection	Subcutaneous Injection
WITHHOLDING PERIOD		
Meat	NIL	NIL
Milk	NIL	NIL
ESI	NIL	NIL

VB12™ is a trademark of Abbey Laboratories Pty Ltd

COBALIFE® is a registered trademark of Bayer AG



VB12™

PLUS SELENIUM FOR SHEEP AND CATTLE

ACTIVE CONSTITUENT

2mg/mL Hydroxocobalamin (as Hydroxocobalamin acetate Vitamin B12), 4mg/mL Selenium as sodium selenate

DIRECTIONS FOR USE

Contraindications

This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.

Precautions

This product should not be administered to sheep with known liver damage concurrently grazing plants which contain pyrrolizidine alkaloids, such as Patterson's Curse or heliotrope.

DOSAGE AND ADMINISTRATION

By subcutaneous injection only.

Species	Dose (mL)
LAMBS	0.5 mL at marking or weaning
SHEEP	1-1.5 mL pre lambing
CALVES	2-3 mL from 2 months of age
COWS	4-6mL pre calving

WITHHOLDING PERIODS

MEAT: Zero (0) days.

MILK: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. Additional information is listed in Safety Data Sheet.

DISPOSAL

Dispose of empty container and packaging by wrapping with paper and putting in garbage.

Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.

STORAGE

Store below 25°C (Air Conditioning). Protect from light.

Store opened product below 25°C.

Use contents within 24 hours of first broaching.

Discard the unused portion.

VB12™ is a trademark of Abbey Laboratories Pty Ltd

APVMA Approval No: 88380/121154