






ZEROMEC™ POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 10g/L IVERMECTIN

Also contains 150g/L Diethylene glycol monobutyl ether (solvent)

Powerful, low volume cattle pour-on

For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle.

-  Low Volume Cattle Pour-On – No Run Off
-  Sustained Activity
-  Kills Internal and External Parasites including Buffalo Flies and Cattle Ticks
-  NIL Milk Withholding Period
-  Pack sizes available: 10L Drum, 2.5L Backpack and 1L Twin Chamber



	ZEROMEC™	BOMECTIN®
Active Constituent	10g/L IVERMECTIN	10g/L IVERMECTIN
Species	Beef and Dairy Cattle	Beef and Dairy Cattle
Dose Rate	1mL/20kg body weight	1mL/20kg body weight
Route of Administration	Pour-on	Pour-on
Withholding periods		
Meat	42 days	42 days
Milk	Zero (0) days	Zero (0) days
ESI	42 days	42 days

ZEROMEC™ is a trademark of Abbey Laboratories Pty Ltd

BOMECTIN® is a registered trademark of Bayer Intellectual Property GmbH

ZEROMETM POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 10g/L IVERMECTIN

Also contains 150g/L Diethylene glycol monobutyl ether (solvent)



For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle

INDICATIONS

For the treatment and control of adult and some immature stages of:

Gastro-intestinal roundworms

Barbers Pole Worm (*Haemonchus placei*)

Hookworm (*Bunostomum phlebotomum*)

Intestinal Threadworm (*Strongyloides papillosus*)

Large Bowel Worm (*Oesophagostomum radiatum*)

Small Intestinal Worm (*Cooperia oncophora*) (*Cooperia punctata*)

Small Brown Stomach Worm (*Ostertagia ostertagi*) (including inhibited stages)

Stomach Hair Worm (*Trichostrongylus axei*)

Thin Necked Intestinal Worm (*Nematodirus spathiger*)

Whipworm (*Trichuris* spp.)

Lungworm (*Dictyocaulus viviparus*)

Sucking and Biting lice

Biting lice (*Bovicola bovis*)

Little sucking lice (*Solenopotes capillatus*)

Long nose sucking lice (*Linognathus vituli*)

Short nosed sucking lice (*Haematopinus eurysternus*)

Mites (*Chorioptes bovis*, *Sarcoptes scabiei* var *bovis*)

Buffalo Fly (*Haematobia irritans exigua*)

Cattle Tick *Rhipicephalus* (*Boophilus*) *microplus*

ZEROMETM POUR-ON FOR CATTLE contains ivermectin, which has a broad spectrum of activity against most internal and external parasites with sensitivity to ivermectin.

Ivermectin is a member of the macrocyclic lactone anthelmintic family of chemicals. Resistance can develop to any chemical.

Maximum tick efficacy is achieved 4-5 days after treatment. In planning a strategic tick control program, it should be noted that viable eggs may have been laid by engorged female ticks that drop from cattle in the initial days after treatment. ZEROMETM POUR-ON FOR CATTLE should be alternated with conventional dips or Pour-Ons when the program requires repeat treatments.

SUSTAINED ACTIVITY

ZEROMETM POUR-ON FOR CATTLE applied at the recommended dose of 1mL per 20kg bodyweight effectively controls infection with *Ostertagia* spp. (Small brown stomach worm) acquired up to 14 days after treatment, with *Dictyocaulus viviparus* (Lungworm) acquired up to 28 days after treatment and *Haematobia irritans exigua* (Buffalo Fly) up to 28 days after treatment.

DIRECTIONS FOR USE

Restraint

Re-treatment interval: DO NOT re-treat animals for 42 days after last treatment.

Side effects

Temporary hair loss occasionally occurs in the application area but will resolve without further treatment.

DOSAGE AND ADMINISTRATION

ZEROMETM POUR-ON FOR CATTLE is for external application to cattle only. The recommended dose rate for ZEROMETM POUR-ON FOR CATTLE is 1mL/20kg bodyweight. Apply along the topline of the animal.

Dose the mob according to the heaviest animal by live weight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing. For animals over 650kg (for example mature bulls) dose at a rate of 1 mL/20kg bodyweight.

Cattle should not be treated with ZEROMETM POUR-ON FOR CATTLE when the hide or hair is wet since the efficacy of the product may be reduced.

Rainfall on the herd within 2 hours of application may result in reduced efficacy of the product.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

MILK: ZERO (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website.

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If products get into the eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

ENVIRONMENTAL PROTECTION

Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used containers.

DISPOSAL

1L: Dispose of empty container by wrapping with paper and putting in garbage.

2.5L, 10L: Triple rinse containers before disposal then dispose of rinsings in a disposal pit away from desirable plants and watercourses.

The disposal of undiluted chemicals is unacceptable. Destroy empty containers by breaking, crushing or puncturing or shredding them when applicable. Dispose of the containers at a local authority landfill readily available in your area. If no landfill is available, bury the containers at a depth of 500mm or more at a licence / approved disposable site. In some states wastes can only be buried at a licenced landfill. Do NOT burn empty containers or product. Clean up spillages promptly.

STORAGE

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval Number: 81646/115688

ZEROMETM is a trademark of Abbey Laboratories Pty Ltd