

QUINDOX™ FEED MEDICATION

ACTIVE CONSTITUENT: 100g/kg OLAQUINDOX

Maximise growth

For use in growing pigs for growth promotion and for the prevention of proliferative enteritis (*campylobacter* spp induced diarrhoea)

- Improvement of liveweight gain and feed conversion
- Easy to use – just mix in feed
- Cost effective



	QUINDOX™	Keyquindox-100™
ACTIVE CONSTITUENT	100g/kg Olaquinox	100g/kg Olaquinox
SPECIES	Pigs	Pigs
ROUTE OF ADMINISTRATION	In feed	In feed
DOSE RATE		
Pigs less than 35kg	1kg / tonne of feed	1kg / tonne of feed
Pigs greater than 35kg	500g / tonne of feed	250-500g / tonne of feed
WITHHOLDING PERIOD	12 hours	12 hours

QUINDOX™ is a trademark of Abbey Laboratories Pty Ltd

Keyquindox-100™ is a registered trademark of International Animal Health Products P/L



QUINDOX™ FEED MEDICATION

ACTIVE CONSTITUENT: 100g/kg OLAQUINDOX

For use in growing pigs for growth promotion and for the prevention of proliferative enteritis (*campylobacter* spp induced diarrhoea)

DIRECTIONS FOR USE

KEEP WHITE PIGS OUT OF DIRECT SUNLIGHT DURING USE.

Thoroughly mix as directed, to ensure even dispersion of the medication in the total ration.

Pigs less than 35kg bodyweight: Add 1kg of QUINDOX™ FEED MEDICATION per tonne of feed.

Pigs greater than 35kg bodyweight: Add 500g QUINDOX™ FEED MEDICATION per tonne of feed.

WITHHOLDING PERIODS

DO NOT ADMINISTER to pigs later than 12 hours before slaughter for human consumption.

SAFETY DIRECTIONS

Harmful if swallowed or will irritate the eyes and may irritate the skin. Avoid contact with eyes and skin and repeated exposure may cause allergic disorders. Do not inhale dust when opening the containers and mixing into feed. Wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and a full face piece respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves and contaminated clothing. If respirator is rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26.

Additional user safety information is listed in the Safety Data Sheet.

DISPOSAL

Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt

STORAGE

Store below 25°C (air conditioning) in a dry place. Store out of direct sunlight. Keep bags tightly closed.

APVMA Approval No. 45799/135166

QUINDOX™ is a trademark of Abbey Laboratories Pty Ltd

May cause an allergic skin reaction.

May cause allergy or asthma symptoms or breathing difficulties if inhaled.

Avoid breathing dust/fume/gas/mist/vapours/spray.

Wear protective gloves/protective clothing/eye protection/face protection.

IF ON SKIN: wash with plenty of soap and water.

IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing.

IF EXPERIENCING RESPIRATORY SYMPTOMS: call a POISON CENTRE or doctor/physician.