


# COXI-QUELL™ 25

## COCCIDIOCIDAL SOLUTION FOR POULTRY



ACTIVE CONSTITUENT: 25g/L TOLTRAZURIL

### Water tight treatment for coccidiosis

COXI-QUELL™ 25 is added to the drinking water of chickens for the treatment and control of coccidiosis caused by Eimeria species including: *Eimeria tenella*, *Eimeria necatrix*, *Eimeria acervulina* and *Eimeria maxima*.


- Can be used as a treatment or preventative for Coccidiosis
- Mixes with all types of water
- Works quickly
- Easy to use
- Cost effective



	COXI-QUELL™ 25	BAYCOX®
ACTIVE	25g/L Toltrazuril	25g/L Toltrazuril
SPECIES	Chickens	Chickens
DILUTION RATE	3L per 1000L water	3L per 1000L water
DOSE RATE	7mg/kg bodyweight/day	7mg/kg bodyweight/day
DOSAGE PERIOD	2 consecutive days for 8 hours	2 consecutive days for 8 hours
WITHHOLDING PERIOD		
Meat	14 days	14 days
Eggs		
Laying Hens	Do not use	Do not use
Replacement Pullets	8 weeks	8 weeks

COXI-QUELL™ is a trademark of Abbey Laboratories Pty Ltd

BAYCOX® is a registered trademark of Bayer Intellectual Property GmbH



# COXI-QUELL™ 25

## COCCIDIOCIDAL SOLUTION FOR POULTRY

ACTIVE CONSTITUENT: 25g/L TOLTRAZURIL

For the treatment and control of coccidiosis in chickens caused by *Eimeria* species.

### INDICATIONS

COXI-QUELL™ 25 is added to the drinking water of chickens for the treatment and control of coccidiosis caused by *Eimeria* species including: *Eimeria tenella*, *Eimeria necatrix*, *Eimeria acervulina* and *Eimeria maxima*.

### DIRECTIONS FOR USE

#### Restraint

DO NOT treat replacement pullets more than twice.

### DOSAGE AND ADMINISTRATION

COXI-QUELL™ 25 is diluted at the rate of 3 litres per 1000 litres of drinking water (3: 1000) and administered on 2 consecutive days with an 8 hour treatment period on both days. Ensure full lighting conditions during treatment. Do not withhold feed during treatment.

Add the required amount of COXI-QUELL™ 25 to treat the birds for 8 hour period and stir thoroughly. Allow for varying water consumption during hot weather.

Treat the birds for 8 hours each day for 2 days (2x8 hour treatments). No other sources of drinking water should be available to birds during the 8 hour treatment period.

### GENERAL DIRECTIONS

DO NOT USE GALVANISED OR RUSTY IRON WATER TANKS AND RETICULATION SYSTEMS.

COXI-QUELL™ 25 mixes with all types of water. However, in areas where the drinking water is acidic, the water should be tested with pH test strips and brought to a minimum pH of 8.5 using sodium carbonate (washing soda).

To avoid excessive foaming, add COXI-QUELL™ 25 to the water after filling the supply tank.

**Broiler Treatment Programme:** To use COXI-QUELL™ 25 in a clinical outbreak obtain an accurate diagnosis. Begin treatment with COXI-QUELL™ 25 as soon as coccidiosis is confirmed. A treatment period of 2 days is sufficient. If required, treatment may be repeated after 5 days.

**Broiler Preventive Programme:** To use COXI-QUELL™ 25 as a coccidiosis preventive in broiler production, give COXI-QUELL™ 25 as above on days 9 and 10, 16 and 17, and 23 and 24 of life. Do not use a coccidiostat in the feed. An additional treatment may be necessary after 35 days of age but not within 14 days of slaughter.

**Layer Birds:** COXI-QUELL™ 25 may be used in replacement pullets and replacement layer breeder birds via the drinking water for an 8 hour period each day for two consecutive days. Treatment may be repeated after 5 days if required.

### WITHHOLDING PERIODS

**MEAT: DO NOT USE** less than 14 days before slaughter for human consumption.

**EGGS: DO NOT USE** in laying hens. **DO NOT USE** in replacement pullets within 8 weeks of laying onset, where eggs are to be used or processed for human consumption. Eggs laid during the withholding period **MUST NOT BE USED** or processed for human consumption.

### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre: Phone Australia 131 126.

### ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

### DISPOSAL

Dispose of container by wrapping with paper and putting in the garbage.

### STORAGE

Store below 30°C (room temperature) in the closed, original container, as cool as possible. Protect from light.

COXI-QUELL™ is a trademark of Abbey Laboratories Pty Ltd

APVMA Approval No:87684/118973