



# EXISHIELD™ GOLD DUAL ACTIVE POUR ON LICE CONTROL

ACTIVE CONSTITUENTS: 35g/L IMIDACLOPRID, 4g/L ABAMECTIN  
Solvent: 300g/L N-METHYL-2-PYRROLIDONE

## Dual Active Pour On Lice Control for Sheep

Fast sheep lice knockdown & up to 5wks protection\*

For the treatment and control of susceptible strains of lice (*Bovicola ovis*) on sheep, including synthetic pyrethroid resistant lice, when applied up to 7 days after shearing. Sheep will be protected against re-infestation with lice for 5 weeks after treatment when applied within 24 hours of shearing.

- ☘ Dual-active lousicides - Cost effective and fast acting
- ☘ Effective immediately on application (shorn sheep within 24 hrs up to 7 days) with up to 5 weeks of protection
- ☘ Easy Pour-On Formulation
- ☘ Bright Blue Colour solution - Treated sheep easily identified
- ☘ Dose rate 50mL per 50kg sheep
- ☘ Kills hatching eggs



	EXISHIELD™ GOLD	Fennec®
Active Constituents	35mg/mL IMIDACLOPRID 4mg/mL ABAMECTIN	35mg/mL IMIDACLOPRID 4mg/mL ABAMECTIN
Species	Sheep	Sheep
Route of Administration	Pour on	Pour on
Dose rate for 50kg Sheep	50mL	50mL
<b>Withholding period</b>		
Meat	63 days	63 days
Milk	Do not use	Do not use
Wool	6 months	6 months
ESI	84 days	84 days

\*On shorn sheep up to 5 days

EXISHIELD™ is a trademark of Abbey Laboratories Pty Ltd

Fennec® is a registered trademark of the Boehringer Ingelheim Group

# EXISHIELD™ GOLD DUAL ACTIVE POUR ON LICE CONTROL

ACTIVE CONSTITUENTS: 35g/L IMIDACLOPRID, 4g/L ABAMECTIN  
Solvent: 300g/L N-METHYL-2-PYRROLIDONE



**For the treatment and control of susceptible strains of lice (*Bovicola ovis*) on sheep, including synthetic pyrethroid resistant lice, when applied up to 7 days after shearing. Sheep will be protected against re-infestation with lice for 5 weeks after treatment when applied within 24 hours of shearing.**

## INDICATIONS

EXISHIELD™ GOLD contains imidacloprid, a member of the neonicotinoid family of chemicals and abamectin, a member of the macrocyclic lactone family of chemicals. Resistance may develop to any chemical.

## DIRECTIONS FOR USE

### Restraint

DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

### Contraindications

DO NOT USE on unshorn sheep, or shorn sheep more than 7 days after shearing. DO NOT USE on ewes with unshorn lambs at foot. This product should not be used on lambs under 6 weeks of age or less than 10kg bodyweight. This product should not be used on severely sick, debilitated or dying animals. This product should not be used on sheep which have not been cleanly shorn for any reason e.g. Mycotic Dermatitis.

### Precautions

This product is for application as a pour on only. Do not give orally or by injection. Do not apply this product at the same time as any other pour on products.

## DOSAGE AND ADMINISTRATION

**Dose rate:** 10mL for each 10kg body weight. Sheep should be weighed before treatment.

Apply the entire dose to sheep under 30kg as a single broadband stripe from poll of the head to the base of the tail.

For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the base of the tail. The stripes should be no further than 10cm apart. Refer to diagrams below.

### Dose table:

Bodyweight (kg)	Dose Volume (mL)	Method	Apply in the pattern below
10 - 12.5	12.5	Single stripe	
12.6 - 15	15		
15.1 - 20	20		
20.1 - 25	25		
25.1 - 30	30		
30.1 - 40	40	Double stripe	
40.1 - 50	50		
50.1 - 60	60		
60.1 - 70	70		
70.1 - 80	80		
80.1 - 90	90		

Animals heavier than 90kg to be dosed at 10mL per 10kg. A representative sample of animals should be weighed before treatment. Dose the mob according to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

## GENERAL DIRECTIONS

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.

Apply product only with the recommended applicator fitted with a six-hole T-bar nozzle. Check the accuracy of equipment before and during treatment. Efficacy is not affected by heavy rainfall before and/or after treatment.

Avoid treating ewes less than 6 weeks before lambing commences, while ewes are lambing or have lambs at foot, as live lice can persist and infect the lambs.

Treated sheep should not be mixed with lice free sheep until 6 weeks after treatment.

Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearing's, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control. EXISHIELD™ GOLD contains a blue dye to identify treated sheep. This dye is scourable from wool. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used. Do not leave container, hose or applicator in the sun for long periods of time. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.

## WITHHOLDING PERIODS

**Meat: DO NOT USE less than 63 days before slaughter for human consumption.**

**Milk: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.**

**Wool: DO NOT USE less than 6 months before shearing or fibre collection.**

**Use of this product may result in wool residues that may not comply with European Union environmental quality standards.**

**Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.**

## TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 84 days before slaughter for export.**

**Before using this product, confirm the current ESI from Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues))**

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes. When using the product wear cotton overalls, buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## ADDITIONAL USER SAFETY INFORMATION

Rehandling Interval: DO NOT handle sheep after treatment until dry.

## ENVIRONMENTAL STATEMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

## DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## STORAGE

Store below 30 °C (Room temperature). Protect from light.

APVMA Approval Number: 92535/138699

EXISHIELD™ is a trademark of Abbey Laboratories Pty Ltd