









TetraMax™ Flexi-Dose INJECTION

ACTIVE CONSTITUENT: 100 mg/mL OXYTETRACYCLINE (AS THE HYDROCHLORIDE)

Kind on the skin and on your pocket

For the treatment of organisms sensitive to oxytetracycline in horses, cattle, sheep, pigs, dogs and cats.

-  Broad spectrum antibiotic
-  Treat multiple species with one product
-  Convenient and flexible dosing
-  Cost effective
-  Pack size: 100mL and 250mL



	TetraMax™ Flexi-Dose	ENGEMYCIN® 100
Active Constituent	100 mg/mL OXYTETRACYCLINE (as the hydrochloride)	100 mg/mL OXYTETRACYCLINE (as the hydrochloride)
Species	Horses, Cattle, Sheep, Pigs, Dogs and Cats	Horses, Cattle, Sheep, Pigs, Dogs and Cats
Withholding Period		
Meat: Cattle & Calves	14 days	14 days
Milk: Cows & Ewes	72 hours (6 milkings)	72 hours (6 milkings)
Meat: Sheep & Pigs	10 days	10 days
Meat: Horses	28 days	28 days
ESI	Not established	Not established

TetraMax™ is a trademark of Abbey Laboratories Pty Ltd.

ENGEMYCIN® is a registered trademark of Intervet International B.V.



TetraMax™ Flexi-Dose INJECTION

ACTIVE CONSTITUENT: 100 mg/mL OXYTETRACYCLINE (AS THE HYDROCHLORIDE)



Kind on the skin and on your pocket

For the treatment of organisms sensitive to oxytetracycline in horses, cattle, sheep, pigs, dogs and cats.

DIRECTIONS FOR USE

Restraints

DO NOT exceed 25mL injection site volume for cattle.

DO NOT exceed 10mL injection site volume for pigs or sheep.

Contraindications

This product is contraindicated in horses during concomitant therapy with corticosteroids.

DOSAGE AND ADMINISTRATION

USE ALL PRODUCT WITHIN 4 WEEKS OF OPENING.

Please refer to leaflet for dose table for each species.

1. 24 hours dose regimen

Give once daily by injection by the route recommended for the species as per the dosing table in leaflet. Care is needed when calculating small dose rates for dogs, cats and young animals. A temporary swelling at the injection site after administration to horses may sometimes be noticed.

2. For long acting mode of action

Injection should be given in the neck region. Give by intramuscular injection in the neck only (cattle, sheep and pigs only), repeat after 2 days, if required.

The maximum recommended dose at any one site is:

Adult cattle: 25mL

Pigs: 10mL

Sheep: 10mL

Injection by all normal parenteral routes. Intravenous injections (horses, cattle and sheep only) must be given slowly over a period of at least one minute. When required, dilution with saline or distilled water is satisfactory, in such cases the solution should be used immediately. Dilution with calcium salts is not recommended as this may lead to precipitation of crystals which makes the solution unsuitable for intravenous use. For long acting dose (cattle, sheep and pigs only) give an intramuscular injection, and repeat after 2 days if required.

TetraMax™ Flexi-Dose can be administered either at a dose rate according to species of 3-8 mg/kg for a 24-hour duration of activity or 10-20 mg/kg for prolonged duration of activity.

Species	BW (kg)	24 HOURS			LONG ACTING		
		Dose (mL)	mg/kg BW	Route	Dosage (mL)	Mg/kg BW	Route
Horses	500	25	5	IM IV	Not recommended		
Adult cattle	500	15	3	IM IV	50	10	IM
Calf	100	8	8	IM IV	20	20	IM
Adult pig	100	5	5	IM	10	10	IM
Piglet	25	2	8	IM	5	20	IM
Sheep	50	4	8	IM IV	10	20	IM
Lamb	15	1.2	8	IM IV	3	20	IM
Dog	25	2	8	IM SC	-	-	-
Cat	5	0.4	8	SC	-	-	-

WITHOLDING PERIODS

MEAT (CATTLE AND CALVES): 14 days.

MEAT (SHEEP AND PIGS): 10 days.

MEAT (HORSES): 28 days.

MILK: 72 hours (cows/ewes).

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI):

This product does not have an ESI established.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Additional information is listed in Safety Data Sheet.

DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE

Store below 25 °C (Air conditioning). Do not freeze. Protect from light. When kept at the recommended storage temperatures, the product remains stable for 14 days after first breaching of the container.

Causes skin irritation.

Causes serious eye irritation.

May cause respiratory irritation.

Do not breathe mist.

Wash contact areas thoroughly after handling.

IF ON SKIN: Wash with plenty of soap and water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

APVMA Approval No: 88734/122118

TetraMax™ is a trademark of Abbey Laboratories Pty Ltd

