

# BEAST™ POUR-ON FOR CATTLE



ACTIVE CONSTITUENT: 87 g/L PERMETHRIN (25:75 cis:trans)

Highly Effective Control against Buffalo Fly and Cattle Lice

BEAST™ POUR-ON FOR CATTLE for the control of buffalo flies\* and control of cattle lice on beef cattle.

Ready-to-use pour on application for cattle

Cost effective

\*Pyrethroid-susceptible strains







	BEAST™	SWAT®
Active Constituent	87g/L Permethrin (25:75 cis:trans)	87g/L Permethrin (25:75 cis:trans)
Species	Cattle	Cattle
Dose rate	1mL/30kg body weight	1mL/30kg body weight
Application	Pour On	Pour On
Pack size	1L, 5L, 10L	1L, 5L
Withholding Period		
Meat: Cattle	7 days	7 days
Milk: Cattle	Do Not Use	Do Not Use
ESI	14 days	14 days

BEAST™ is a trademark of Abbey Laboratories Pty Ltd

SWAT® is a trademark of Pharmachem Australia Pty Ltd

# BEAST™ POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 87 g/L PERMETHRIN (25:75 cis:trans)







### For control of buffalo flies\* and control of cattle lice on beef cattle

Pyrethroid-susceptible strains

## DIRECTIONS FOR USE RESTRAINTS

DO NOT USE on lactating or pregnant cows where milk or milk products may be used or processed for human consumption.

DO NOT TREAT animals with this product more than 3 times in any given 12 month period.

Re-treatment interval: DO NOT RE-TREAT animals for 21 days after last treatment.

More frequent use of this product is likely to result is residues above the MRL.

#### **PRECAUTIONS**

Use with caution as the safety of this product in calves has not been established.

### DOSAGE AND ADMINISTRATION Shake well before use.

Animal & Target Pest	Dosage	Administration
Cattle, Ready-to-use Pour-On Application: Buffalo Flies (Haematobia irritans exigua) Cattle Lice, biting and sucking species (Bovicola bovis, Linognathus vituli, haematopinus eurysternus)	1 mL/30 kg body weight per animal up to 15 mL maximum. Reapply as needed but not more often than once every 21 days.	Apply by pouring from the calibrated chamber of the dispenser.  For Buffalo Fly control: Apply along the backline, from the shoulders to the tail butt, when fly number exceed acceptable levels.  For Cattle Lice control: Apply along the backline from the head poll down the neck and over the shoulders. For optimum residual control, reapply after 3 weeks.

**Application:** Loosen cap and squeeze bottle to fill measuring chamber with required dosage. Remove cap and pour along the animal's backline. Apply 1mL/30kg body weight up to a maximum of 15 mL per animal. The maximum number of applications is 3 per season.

#### **GENERAL DIRECTIONS**

BEAST™ POUR-ON FOR CATTLE will give effective control of Buffalo Flies and Cattle Lice when used as directed. The level and duration of control may be reduced due to high fly pressure. Treat all animals in the group.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

#### WITHHOLDING PERIODS

MEAT: DO NOT USE less that 7 days before slaughter for human consumption.

#### TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 14 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal health Pty Ltd on 02 8088 0720 or the APVMA website (www.apvma.gov.au/residues).

#### **SAFETY DIRECTIONS**

May irritate the eyes. Avoid contact with eyes. Wash hands after use.

#### **FIRST AID INSTRUCTIONS**

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131 126.

#### **ENVIRONMENT PROTECTION**

This product is highly toxic to fish. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

#### **DISPOSAL**

#### 1 L containers:

Dispose of container by wrapping with paper and putting in garbage.

#### 5 L and 10 L containers:

Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### **STORAGE**

Store below 30 °C (room temperature). Store upright in the original container tightly closed, in a dry place. DO NOT store in direct sunlight.

APVMA Approval Number: 92938/137389



