

# FLUZOLE™ Oral Wormer for Pigs and Poultry

ACTIVE CONSTITUENT: 50 mg/g FLUBENDAZOLE

## No Worms. No Worries.

For use as an aid and control of sensitive strains of internal parasites in pigs and poultry.

- ✿ Easy to use – just mix into feed!
- ✿ Effective against all stages of worms in poultry including tapeworm
- ✿ Zero day WHP for eggs
- ✿ Zero effect on:
  - egg production, fertility or hatchability in chickens<sup>1</sup>
  - fertility or embryo formation in pigs<sup>2</sup>




	FLUZOLE™	Flubenol™
Active Constituent	50mg/g Flubendazole	50mg/g Flubendazole
Species	Pigs & Poultry	Pigs & Poultry
Application	In Feed	In Feed
<b>Dose Rates</b>		
Poultry	30g for 7 consecutive days	30g for 7 consecutive days
Poultry incl. Tapeworm	60g for 7 consecutive days	60g for 7 consecutive days
Pigs	30g for 10 consecutive days	30g for 10 consecutive days
<b>Withholding Periods</b>		
Meat: Chicken	7 days	7 days
Meat: Pigs	7 days	7 days
Eggs	Zero (0) days	Zero (0) days
ESI	7 days	7 days

FLUZOLE™ is a trademark of Abbey Laboratories Pty Ltd

Flubenol™ is a trademark of Elanco Australasia Pty Ltd

# FLUZOLE™ Oral Wormer for Pigs and Poultry

ACTIVE CONSTITUENT: 50 mg/g FLUBENDAZOLE



## For use as an aid and control of sensitive strains of internal parasites in pigs and poultry

### INDICATIONS

FLUZOLE™ contains Flubendazole, a member of the benzimidazole group family of chemicals.

FLUZOLE™ can be used as an aid for the control of sensitive strains of the following internal parasites: Gastrointestinal worms (*Ascaridia galli*, *Heterakis gallinarum*, *Capillaria spp.*, *Raillietina spp.*) in chickens and gastrointestinal worms (*Ascaris suum*, *Hyostrongylus rubidus*, *Oesophagostomum dentatum*) and lungworms (*Metastrongylus apri*) in pigs. Resistance may develop to any chemical.

### DIRECTIONS FOR USE

#### Contraindication

This product is contraindicated for use in pigeons and parrots.

#### Precaution

Do not overdose.

### DOSAGE AND ADMINISTRATION

Use treated feeds within 6 months of preparation.

#### CHICKENS

*Ascaridia galli*, *Heterakis gallinarum*, *Capillaria spp.*

600g of FLUZOLE™ per tonne of feed (equivalent to 30g Flubendazole (30 ppm)) for 7 consecutive days.

For small numbers of animals, 13g of FLUZOLE™ is mixed into 21.5kg of compound feed for chickens.

#### TAPEWORM

*Raillietina spp.*

1200g of FLUZOLE™ per tonne of feed (equivalent to 60g Flubendazole (60 ppm)) for 7 consecutive days.

Ensure birds have access to feed to enable consumption of the required dose. Optimum results are achieved when using stringent chicken housing hygiene requirements.

### PIGS

*Ascaris suum*, *Hyostrongylus rubidis*, *Oesophagostomum dentatum* and *Metastrongylus apri*

600g of FLUZOLE™ per tonne of feed (equivalent to 30g Flubendazole (30 ppm)) for 10 consecutive days.

#### Individual and single administration to breeding pigs

FLUZOLE™ is mixed into the feed at a dose of 5mg Flubendazole per kg body weight. This corresponds to 13g of FLUZOLE™ per 130kg of body weight.

FLUZOLE™ should be thoroughly mixed into feed to ensure even distribution of the medication. Feed can be administered as either a mash, crumble or pellet.

### WITHHOLDING PERIODS

**MEAT (CHICKEN): REMOVE ALL MEDICATED FEED 7 days before slaughter for human consumption.**

**EGGS: Zero (0) days.**

**MEAT (PIGS): REMOVE ALL MEDICATED FEED 7 days before slaughter for human consumption.**

### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Laboratories Pty Ltd on 02 8088 0720 or the APVMA website ([apvma.gov.au/residues](http://apvma.gov.au/residues)).**

### SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131 126. Additional information is listed in the safety data sheet (SDS).

### ENVIRONMENT PROTECTION

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

### DISPOSAL

Single-rinse or shake container into the medicated feed. Do not dispose of undiluted chemicals on-site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### STORAGE

Store below 30°C (room temperature). Use treated feed within 6 months of preparation.

### APVMA Approval Number: 93315/138712

<sup>1</sup> Froyman R. & De Keyser H. Flubendazole: safety regarding egg production and reproductive performance of breeder chickens. Av Dis 27:1 (43-48). 1983.

<sup>2</sup> Committee for Medicinal Products for Veterinary Use. Flubendazole (extrapolation to poultry). Summary Report (4). European Medicines Agency. July 2006.

FLUZOLE™ is a trademark of Abbey Laboratories Pty Ltd