







LEVATAK™ Soluble Powder

ACTIVE CONSTITUENT: 850mg/g LEVAMISOLE (as Levamisole Hydrochloride)

In Water Wormer

For the control of Levamisole susceptible gastrointestinal roundworms of domestic poultry, pigeons, aviary and pet birds, and birds in zoological collections.

For use in treatment of roundworm (*Ascaridia* spp.), hairworm (*Capillaria* spp.), caecal worm (*Heterakis* spp.).

-  Broad spectrum control
-  Easy to use – just mix into water!
-  Zero day egg WHP
-  Cost effective



	LEVATAK™ Soluble	CCD® Levamisole
Active Constituent	850mg/g LEVAMISOLE (as Levamisole Hydrochloride)	850mg/g LEVAMISOLE (as Levamisole Hydrochloride)
Species	Poultry	Poultry
Administration method	Oral drinking water	Oral drinking water
Dose rate	28mg/kg live weight	28mg/kg live weight
Withholding Period		
Meat	7 days	7 days
Eggs	Zero (0) days	Zero (0) days

LEVATAK™ is a trademark of Abbey Laboratories Pty Ltd

CCD® is a registered trademark of CCD Animal Health Pty Ltd

LEVATAK™ Soluble Powder

ACTIVE CONSTITUENT: 850mg/g LEVAMISOLE (as Levamisole Hydrochloride)



For the control of Levamisole susceptible gastrointestinal roundworms of domestic poultry, pigeons, aviary and pet birds, and birds in zoological collections. For use in treatment of roundworm (*Ascaridia* spp.), hairworm (*Capillaria* spp.), caecal worm (*Heterakis* spp.).

DIRECTIONS FOR USE

CONTRAINDICATIONS

Do not use in extremely hot weather. Not for use in species or strains of birds to be sensitive to the effects of Levamisole, and not for use in birds that are feeding young. Parrots appear to be less tolerant than pigeons to levamisole.

DOSAGE AND ADMINISTRATION

28mg/kg live weight or as directed by a veterinary surgeon. As a guide, assuming medicated water intake of 35mL/bird during the treatment period, use 800g LEVATAK™ per 900-1000L drinking water, or 8g per 10L water for small number of birds. The amount of solution prepared should be as much as will be consumed in twelve hours. Remove other sources of medicated water.

WITHHOLDING PERIODS

EGGS: Zero (0) days.

MEAT: REMOVE ALL MEDICATED WATER 7 days before slaughter for human consumption.

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the eyes and skin. May irritate the nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin.

When opening the container and preparing product for use and mixing, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves, goggles or safety glasses and disposable dust mask covering mouth and nose. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water.

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles or safety glasses and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Additional information is listed in Safety Data Sheet.

DISPOSAL

Single-rinse or shake container into the medicated water. Do not dispose of undiluted chemicals on-site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30 °C (Room temperature) in a dry place. Protect from light. Keep container tightly closed.

**APVMA Approval Number:
88799/122348**