




BOVIMEC™ POUR-ON FOR BEEF AND DAIRY CATTLE

ACTIVE CONSTITUENT: 5g/L IVERMECTIN

Effective parasite protection and coverage for beef and dairy cattle

For the treatment and control of Ivermectin-sensitive internal and external parasites of beef and dairy cattle.

- 🌿 Pour On convenience with high volume beast coverage
- 🌿 Broad parasite cover - Kills internal and external parasites including Buffalo Flies and Cattle Ticks
- 🌿 Milk Safe – NIL Withholding Period
- 🌿 Wide safety margin
- 🌿 Rain Safe
- 🌿 Dung beetle & earthworm low impact
- 🌿 Blue coloured solution – Ease of application identification
- 🌿 Cost effective choice
- 🌿 Pack sizes: 1L Twin Chamber Bottle, 5L Backpack, 20L Drum



	BOVIMEC™ POUR-ON	IVOMECS® POUR-ON
Active Constituent	5g/mL	5g/mL
Species	Cattle - Beef and Dairy	Cattle - Beef and Dairy
Dose rate	1mL/10kg	1mL/10kg
Route of Administration	Pour On	Pour On
Withholding Period		
Meat	28 days	42 days
Milk	Nil	28 days
Export Slaughter Interval	28 days	42 days

BOVIMEC™ is a trademark of Abbey Laboratories Pty Ltd

IVOMECS® is a registered trademark of Bayer Intellectual Property GmbH

BOVIMEC™ POUR-ON FOR BEEF AND DAIRY CATTLE

ACTIVE CONSTITUENT: 5g/L IVERMECTIN



For the treatment and control of Ivermectin-sensitive internal and external parasites of beef and dairy cattle.

INDICATIONS

For the treatment and control of Ivermectin sensitive:

Gastrointestinal nematodes:

- Adult and Immature (L4): Small Brown Stomach Worms (including inhibited stages) (*Ostertagia ostertagi*), Barber's Pole Worm (*Haemonchus placei*), Stomach Hair Worm (*Trichostrongylus axei*), Intestinal Hair Worm (*T. colubriformis*), Small Intestinal Worm (*Cooperia spp.*), Nodule Worm (*Oesophagostomum radiatum*); and
- Adult stages of *Cooperia oncophora*, *C. punctata*, *Oesophagostomum venulosum*, Whip Worm (*Trichuris spp.*), Intestinal Threadworm (*Strongyloides papillosis*); and
- Immature stages (L4) of Thin Necked Intestinal Worm (*Nematodirus spp.*)

Lungworms: (adult and immature) *Dictyocaulus viviparus*.

Eyeworm: *Thelazia spp.* (adult stages).

Sucking and Biting Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Bovicola bovis*, *Solenopotes capillatus*.

Mites: *Chorioptes bovis*, *Sarcoptes scabiei var bovis*.

Buffalo Flies: *Haematobia irritans exigua*.

Cattle Ticks: (*Boophilus microplus*) including organophosphate, synthetic pyrethroid and amidine-resistant strains.

PERSISTENT ACTIVITY

BOVIMEC™ POUR-ON applied at the recommended dose of 1mL per 10kg liveweight controls infections of Stomach Hair Worm (*Trichostrongylus axei*) and Small Intestinal Worm (*Cooperia spp.*) for up to 14 days after treatment; Small Brown Stomach Worm (*Ostertagia spp.*) and Nodule Worm (*Oesophagostomum radiatum*) for up to 21 days after treatment; Lungworms (*Dictyocaulus viviparus*) for up to 28 days after treatment; Buffalo Fly (*Haematobia irritans exigua*) for up to 21 days after treatment.

Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control programme. BOVIMEC™ POUR-ON should be alternated with conventional dips or pour-ons when the programme requires repeated treatments.

For best results BOVIMEC™ POUR-ON should be part of a total parasite control programme including internal and external parasites. Consult your veterinarian for a complete parasite control programme.

DIRECTIONS FOR USE

Restraint

DO NOT USE in pregnant cows less than 7 days before calving.

DOSAGE AND ADMINISTRATION

This product is formulated only for external application to cattle; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tail head. The dose is 1mL for each 10kg of liveweight.

Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc.). When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid overdosing. Do not underdose. Cattle should be weighed with either scales or with a weighband.

Liveweight (kg)	Dose (mL)	1L Twin Chamber	5L Backpack	20L Drum
<100	1mL/10kg	≥100	≥500	≥2000
101-150	15	66	333	1333
151-200	20	50	250	1000
201-250	25	40	200	800
251-300	30	33	166	666
301-350	35	28	142	571
351-400	40	25	125	500
401-450	45	22	111	444
451-500	50	20	100	400
501-550	55	18	90	363
551-600	60	16	83	333
601-650	65	15	76	307

Heavier animals (for example, mature bulls) should receive an additional 5mL for each 50 kg over 650 kg. Treatment should not be repeated in less than 42 days.

Administration

Efficacy of BOVIMEC™ POUR-ON is not adversely affected when the hide is wet or if rainfall occurs shortly after treatment. However, it is not considered good agricultural practice to treat animals if they are wet or rain is likely.

GENERAL DIRECTIONS

NOTE TO USER

The colour of BOVIMEC™ POUR-ON may fade when exposed to light. This loss of colour does not reflect loss of potency of ivermectin. However, prolonged exposure (i.e. weeks) to light can result in a gradual decline of ivermectin potency in the formulation.

SAFETY TO CATTLE

Field studies have demonstrated a wide safety margin. The topically applied formulation is expected to be well tolerated by breeding animals.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: ZERO (0) days.

Bobby calves: DO NOT SLAUGHTER for 4 days bobby calves born to cows treated during pregnancy.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL

(ESI): DO NOT USE less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval.

Before using this product confirm the current ESI from Abbey Animal Health Pty Ltd on (02) 8088 0720 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. If product gets on skin, immediately wash the area with soap and water. If product gets in eyes, wash out immediately with water. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

For further information, refer to the Safety Data Sheet.

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

When used according to label directions BOVIMEC™ POUR-ON is not likely to have an unintended effect that is harmful to the environment or have an overall adverse effect on dung beetle or earthworm populations. Increased mortality and impaired development of dung beetle larvae may be expected, but for a limited period after treatment. There may also be short-term indirect effect on the rate of earthworm colonisation of dung pats.

DISPOSAL

Triple or (preferably) pressure rinse container. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to a recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured container 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30 °C (room temperature). Protect from light.

APVMA Approval Number: 82468/105807

BOVIMEC™ is a trademark of Abbey Laboratories Pty Ltd