



PARAFEN™ 100 ORAL ANTHELMINTIC FOR CATTLE AND HORSES

ACTIVE CONSTITUENT: 100g/L FENBENDAZOLE

Effective treatment & control of parasites in cattle and horses

For control of benzimidazole sensitive mature and immature roundworms and lungworm in cattle, and for the control of large strongyles, small strongyles, ascarids and pinworms in horses.

- 🌿 Broad spectrum oral drench for Cattle and Horses
- 🌿 Ready to use, no dilution or mixing is necessary
- 🌿 Effective against Barbers Pole worm, Small brown stomach worm, stomach Hair Worm, Small Intestinal Worm, Thin Necked Intestinal Worm, Nodule worm, Whipworm, Capillaria spp., Lungworms, Hookworm in cattle
- 🌿 For control of Large strongyles, Small strongyles, Ascarids, Pinworms in horses. Aids in control of tapeworm.
- 🌿 Safe for pregnant mares and working stallions
- 🌿 Dairy Cattle safe - Nil Milk Withholding Period
- 🌿 Cost effective
- 🌿 Pack sizes: 1L Backpack, 20L Drum



	PARAFEN™ 100	PANACUR® 100
Active Constituent	100g/L FENBENDAZOLE	100g/L FENBENDAZOLE
Species	Cattle & Horses	Cattle & Horses
Dose Rate		
Cattle	3.75mL per 50kg bodyweight	3.75mL per 50kg bodyweight
Horses	1mL per 10kg bodyweight	1mL per 10kg bodyweight
Foals (to control Strongyloides westeri)	5mL per 10kg bodyweight	5mL per 10kg bodyweight
Route of Administration	Oral Drench	Oral Drench
Withholding Period		
Meat (Cattle)	14 days	14 days
Milk (Cattle)	Nil	Nil
ESI (Cattle)	14 days	14 days
Meat (Horses)	28 days	28 days

PARAFEN™ is a trademark of Abbey Laboratories Pty Ltd

PANACUR® is a registered trademark of Coopers Animal Health (A division of Intervet Australia Pty Ltd)

PARAFEN™ 100 ORAL ANTHELMINTIC FOR CATTLE AND HORSES

ACTIVE CONSTITUENT: 100g/L FENBENDAZOLE



Effective treatment & control of parasites in cattle and horses

INDICATIONS

PARAFEN™ 100 contains fenbendazole, a member of the benzimidazole family of drenches. PARAFEN™ 100 is active against both mature (adult) and immature (larval) fenbendazole sensitive nematode parasites. PARAFEN™ 100 has activity (80-90%) against inhibited (hypobiotic) larvae of *Ostertagia* in cattle. PARAFEN™ 100 is effective against sensitive strains of the following internal parasites:

CATTLE: Barbers Pole worm (*Haemonchus spp.*), Small brown stomach worm (*Ostertagia spp.*), Stomach Hair Worm (*Trichostrongylus spp.*), Small Intestinal Worm (*Cooperia spp.*), Thin Necked Intestinal Worm (*Nematodirus spp.*), Nodule worm (*Oesophagostomum Radiatum spp.*), Whipworm (*Trichuris spp.*), *Capillaria spp.*, Lungworms (*Dictyocaulus Viviparus*), Hookworm (*Bunostomum Phlebotomum*). PARAFEN™ 100 also aids in the control of Tapeworm (*Moniezia spp.*).

HORSES: Large strongyles (*Strongylus vulgaris*, *S.edentatus*, *S.equinus*), Small strongyles (*Cyathostomum spp.*, *Tridontophorus spp.*), Ascarids (*Parascaris Equorum*), Pinworms (*Oxyuris Equi*).

Resistance may develop to any chemical.

DIRECTIONS FOR USE

Contraindications

This product should not be used in dogs or any other pets. This product is not recommended for use in goats as safety and efficacy have not been evaluated.

DOSAGE AND ADMINISTRATION

Shake container before use. PARAFEN™ 100 is ready to use, no dilution or mixing is necessary. PARAFEN™ 100 may be administered orally through standard drenching equipment. After use, wash equipment in clean water.

CATTLE

Dose Rate: 3.75mL PARAFEN™ 100 per 50kg bodyweight (7.5mg fenbendazole/kg bodyweight).

For animals over 650kg dose rate of 3.75 mL per 50kg bodyweight.

A representative sample of animals should be weighed before treatment either with scales or weigh band. Dose according to the heaviest animal by live weight in the group (cows, bulls, heifers, etc). Do not underdose where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.

Dose Rate Guide:

Body weight (kg)	Dose Volume (mL)	Body weight (kg)	Dose Volume (mL)
Up to 50	3.75	351-400	30.0
51-100	7.5	401-450	33.75
101-150	11.25	451-500	37.5
151-200	15.0	501-550	41.25
201-250	18.75	551-600	45.0
251-300	22.5	601-650	48.75
301-350	26.25		

HORSES

PARAFEN™ 100 can be administered by dosing syringe or drenching gun. PARAFEN™ 100 has a safety margin in excess of 50 times the recommended dose. PARAFEN™ 100 may be safely administered to pregnant mares and working stallions.

Dose Rate: 1mL PARAFEN™ 100 per 10 kg body weight (10mg fenbendazole/kg body weight). To control *Strongyloides westeri* in foals a dose of 5mL per 10 kg (50mg fenbendazole/kg body weight) is required.

For animals over 500 kg dose rate of 1mL per 10kg bodyweight.

Dose Rate Guide:

Body weight (kg)	Dose Volume (mL)
Up to 50	5.0
51-100	10.0
101-150	15.0
151-200	20.0
201-250	25.0
251-300	30.0
301-350	35.0
351-400	40.0
401-450	45.0
451-500	50.0

When used as directed no specific precautions are required other than those normally associated with the careful handling of stock especially weak or pregnant animals. Internal parasites may develop resistance to drenches (anthelmintics). If drench failure with this preparation is suspected, seek veterinary confirmation and notify manufacturer.

GENERAL DIRECTIONS

PARAFEN™ 100 does not stain or predispose animals to photosensitization. PARAFEN™ 100 is palatable and readily accepted by livestock. PARAFEN™ 100 may be administered concurrently with vaccines, pour-on lice treatments and dipping operations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT (Cattle): DO NOT USE less than 14 days before slaughter for human consumption.

MILK: ZERO (0) days.

MEAT (Horses): DO NOT USE less than 28 days before slaughter for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI) -

CATTLE: DO NOT USE less than 14 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website (apvma.gov.au/residues).

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26.

Additional user safety information is listed in the Safety Data Sheet.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature). Do not freeze. Store in tightly closed original container in a well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval Number: 92976/137485

Harmful if swallowed.

Harmful in contact with skin.

Do not get in eyes, on skin, or on clothing.

Wash contacted areas thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wash contaminated clothing before use.

IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water

PARAFEN™ is a trademark of Abbey Laboratories Pty Ltd