





RWR MOR-N-MECTIN Oral Horse Worming Paste

ACTIVE CONSTITUENTS: 167mg/g MORANTEL TARTRATE, 4mg/g ABAMECTIN

Comprehensive worming in one syringe

For the treatment and control of tapeworms (*A. perfoliata*), roundworms, (including arterial larval stages of *Strongylus vulgaris* and benzimidazole resistant small strongyles), bots and skin lesions caused by *Habronema* spp. and *Draschia* spp. (summer sores) and *Onchocerca* spp. *microfilariae* (cutaneous onchocerciasis) in horses.

-  Dual active ingredients
-  Broad spectrum coverage
-  Safe to use in pregnant mares, stallions and foals from 4 weeks of age
-  Convenient dial-a-dose syringe



	RWR MOR-N-MECTIN	Ammo® Allwormer
Active Constituents	167mg/g MORANTEL TARTRATE, 4mg/g ABAMECTIN	167mg/g MORANTEL TARTRATE, 4mg/g ABAMECTIN
Species	Horses	Horses
Route of Administration	Oral	Oral
Pack Size	32.5g	32.5g
Dose Rate	5mL/100kg bodyweight	5mL/100kg bodyweight
Australian Owned Company	Yes	No

AMMO® is a registered trademark of Ceva Animal Health Pty Ltd

RWR MOR-N-MECTIN Oral Horse Worming Paste

ACTIVE CONSTITUENTS: 167mg/g MORANTEL TARTRATE, 4mg/g ABAMECTIN



For the treatment and control of tapeworms (*A. perfoliata*), roundworms, (including arterial larval stages of *Strongylus vulgaris* and benzimidazole resistant small strongyles), bots and skin lesions caused by *Habronema* spp. and *Draschia* spp. (summer sores) and *Onchocerca* spp. *microfilariae* (cutaneous onchocerciasis) in horses.

INDICATIONS

At the dosage rate of 5.4g per 100kg bodyweight RWR MOR-N-MECTIN is indicated for the treatment and control of the following parasites in horses:-

Ascarids: *Parascaris equorum* (adult and immature)

Bots: *Gasterophilus* spp. (oral and gastric stages)

Hairworms: *Trichostrongylus axei* (adult)

Intestinal Threadworms: *Strongyloides westeri* (adult)

Large Mouth Stomach Worms:

Habronema muscae (adult)

Large and Small Strongyles: *Strongylus vulgaris* (adults and arterial larval stages), *Strongylus edentatus* (adult and tissue stages), *Strongylus equinus* (adults) and *Triodontophorus* spp. (adults). *Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Gyalocephalus* spp.

Lungworms: *Dictyocaulus arnfieldi* (adult and immature)

Neck Threadworms: *Onchocerca* spp. (*microfilariae*)

Pinworms: *Oxyuris equi* (adult and immature)

Tapeworms: *Anoplocephala perfoliata*

RWR MOR-N-MECTIN also controls skin lesions caused by *Habronema* and *Draschia* spp., cutaneous larvae (summer sores), and *Onchocerca* spp. *microfilariae* (cutaneous onchocerciasis).

RWR MOR-N-MECTIN is highly effective in the treatment and control of most types of equine helminths including tapeworms, gastrointestinal, cutaneous and pulmonary nematodes and bots of horses of any age. With a regular treatment program the chances of verminous arteritis and colic caused by *S. vulgaris* will be reduced.

Consult a veterinarian when developing a worm control program to suit your needs. Resistance may develop to any chemical.

DIRECTIONS FOR USE

DOSAGE AND ADMINISTRATION

Given orally at the recommended dose rate of 5.4g or 5mL/100kg bodyweight. One syringe is capable of treating horses of weight up to 600kg. The dose of 5.4g/100kg bodyweight delivers 0.22mg/kg of ABAMECTIN and 9.0mg/kg of MORANTEL TARTRATE.

Screw the plastic adjuster out to the weight of the horse, insert the syringe into the side of the horse's mouth between the front and back teeth and deposit the paste towards the back of the tongue by depressing the plunger quickly as far as it will go. Immediately raise the horses' head for a few seconds to ensure the full dose has been accepted.

General Directions

Develop a regular treatment program: All horses should be subject to regular worming to maintain good health and peak condition. Particular attention should be given to young horses and mares. Foals should be treated initially at 6 to 8 weeks of age and this treatment repeated regularly as required.

RWR MOR-N-MECTIN displays a wide safety margin at the recommended dose rate of 5.4g/100kg bodyweight and may be used in both young foals to very old animals. Mares may be treated at any stage of pregnancy and fertility will not be affected in either sex.

WITHHOLDING PERIODS

MEAT WITHHOLDING PERIODS (HORSES):

DO NOT USE less than 28 days before slaughter for human consumption.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

ENVIRONMENTAL PROTECTION

Abamectin is extremely toxic to aquatic life. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of container by wrapping with paper and putting in garbage.

STORAGE

Store below 30° C (air conditioning). Protect from light.

APVMA Approval Number: 69678/137517