

Abbey™ Flunixx PASTE

ACTIVE CONSTITUENT: 50mg/g FLUNIXIN (equivalent to 83mg/g Flunixin meglumine)

Rapid, targeted relief with easy administration for horse owners

For the alleviation of inflammation and pain associated with musculoskeletal disorders in horses.

- ❁ Potent NSAID with analgesic, anti-inflammatory and antipyretic action
- ❁ Rapidly absorbed after oral administration, with 80% bioavailability
- ❁ Onset of therapeutic effect usually occurs within 2 hours
- ❁ Peak response within 15 hours and duration of effect up to 30 hours



Abbey™ Flunixx Paste

Active Constituent	50mg/g FLUNIXIN (equivalent to 83mg/g Flunixin meglumine)
Species	Horses
Dose rate	1.1mg/kg bodyweight once daily for up to 5 days
Route of Administration	Oral
Australian Made	Yes
Australian Owned	Yes

ABBEY™ is a trademark of Abbey Laboratories Pty Ltd

Abbey™ Flunixx PASTE

ACTIVE CONSTITUENT: 50mg/g FLUNIXIN (equivalent to 83mg/g Flunixin meglumine)



For the alleviation of inflammation and pain associated with musculoskeletal disorders in horses.

INDICATIONS

For use by or under direction of a veterinarian.

Mode of Action

The active constituent in Abbey™ Flunixx Paste is a potent non-steroidal, non-narcotic analgesic with anti-inflammatory, antipyretic activity.

DIRECTIONS FOR USE

Use as directed by prescribing veterinarian.

Contraindications

Not to be used in animals still under the effects of general anaesthesia.

Not to be used in animals suffering from cardiac, hepatic or renal disease, where there is possibility of gastrointestinal ulceration or bleeding, or where there is evidence of blood dyscrasia or hypersensitivity to the product.

Not to be used concurrently with any animal remedy that is potentially nephrotoxic, especially if the animal is dehydrated or hypotensive.

Do not exceed recommended dose rate or duration of treatment.

Precautions

The effect of Abbey™ Flunixx Paste on pregnancy has not been determined.

Use with caution when administering to aged horses and foals, especially when under 6 weeks of age.

Monitor drug capability when any other concomitant therapy is needed. Include appropriate antimicrobial therapy when treating inflammatory conditions that are likely to be associated with bacterial infection.

Side effects

Gastric irritation and ulceration may occur if dose rate and/or duration of treatment is increased over recommended levels.

DOSAGE AND ADMINISTRATION

For oral administration only.

The recommended dose of Flunixin is 1.1mg/kg bodyweight once daily for up to 5 days.

Turn the syringe dial to the appropriate mark to deliver the correct dose. Each syringe is sufficient to treat one 455kg horse once daily for 3 days.

Insert the nozzle of the syringe carefully through the interdental space and deposit the required amount of paste on the back of the tongue by depressing the plunger. Raise the horse's head for few seconds to ensure the dose has been accepted.

GENERAL DIRECTIONS

If used in performance animals, the regulations of the relevant authorities regarding medication should be observed.

WITHHOLDING PERIODS

MEAT (HORSES): DO NOT USE less than 28 days before slaughter for human consumption.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

DISPOSAL

Dispose of container by wrapping with paper and putting in garbage.

STORAGE

Store below 30°C (room temperature).

APVMA Approval Number: 95770/147110