

EXIFLEA® PLUS

3-in-1 DEFENCE

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

This leaflet is for:

- Puppies & Small Dogs up to 4kg
- Dogs 4-10kg
- Dogs 10-25kg
- Dogs over 25kg

EXIFLEA® PLUS 3-in-1 Defence

ACTIVE CONSTITUENTS:
100g/L IMIDACLOPRID, 25g/L MOXIDECTIN

Fleas, Heartworms and Worms:

- Once a month topical flea and worm treatment
- Stops fleas feeding in 3-5 minutes
- Prevents heartworm and controls intestinal worms & larvae

INDICATIONS

For the treatment and prevention of fleas *Ctenocephalides* spp. Reduces the incidence of Flea allergy dermatitis. Larval flea stages are killed in the surroundings of EXIFLEA® PLUS treated pets. For the prevention of heartworm *Dirofilaria immitis* infections. One application lasts for a month in dogs.

For the treatment and control of:

Roundworms

Toxocara canis, *Toxascaris leonina* kills Larval (L4), immature and adult worms

Hookworms

Ancylostoma caninum, *Ancylostoma braziliense* (this hookworm is usually restricted to tropical areas in Australia), *Ancylostoma ceylanicum*, *Uncinaria stenocephala*, kills larval (L4), immature and adult worms of all species.

Whipworm

Trichuris vulpis

Generalised demodicosis

Demodex canis

Sarcoptic mange

Sarcoptes scabiei

Ear mites

Otodectes cynotis

Lice

Trichodectes canis, *Linognathus setosus* for up to 6 weeks

DIRECTIONS FOR USE

Consult a veterinarian before use. Treatment for heartworm should occur regularly at monthly intervals. If worm problem persists, consult a veterinarian.

For external use only.

DO NOT GIVE PRODUCT BY MOUTH.

Precautions

DO NOT apply to cats.

Use with caution in sick, debilitated or underweight animals. EXIFLEA® PLUS may be safely applied to heartworm infected dogs therefore there is no absolute requirement for a heartworm test before commencing the treatment. However, a diagnostic test for heartworm prior to commencement of treatment is recommended to ensure the optimum health of dogs.

EXIFLEA® PLUS has no strong chemical smell. The liquid vehicle in EXIFLEA® PLUS may discolour or damage certain surfaces and fabrics e.g. polished floorboards, leather. Exclude the animal from contact with such surfaces until the product dries on animal's coat.


DOSAGE AND ADMINISTRATION

Dosage table

For animals of this type	Use 1 tube from this pack colour
Puppies & small dogs up to 4kg	Green
Dogs 4-10kg	Turquoise
Dogs 10-25kg	Maroon
Dogs 25-40kg	Blue

For very large dogs over 40kg, apply 2 tubes of EXIFLEA® PLUS for Dogs over 25kg (blue) by dividing the dose between the shoulder blades, on the midline of the back between hips, and at one point in between.

- 1 Remove required tube(s) from the package.



- 2 Hold tube upright and pull off cap.
- 3 Reverse the cap and push cap back on tube.
- 4 Twist cap to break the seal, remove cap from tube.
- 5 Site of application

DOGS UNDER 25kg	DOGS 25-40kg
1 TUBE Apply between the shoulder blades	1 TUBE Apply by dividing the dose between the shoulder blades, on the midline of the back between hips, and at one point in between blades

- 6 The animal should be standing for easy application. Remove any collar prior to application and replace when product is dry. Part the hair until the skin is visible. Place the tip of the tube on the skin and squeeze the tube firmly twice to apply the solution directly on the skin. Avoid pet's eyes OR mouth.
- 7 Record treatment date and repeat monthly or as required. EXIFLEA® PLUS remains efficacious following shampoo treatment, swimming or exposure to rain. If washing animal during use of this product use a soap free shampoo.
- 8 Occasionally hair loss may occur at the application site. This is usually a single occurrence and hair will grow back uneventfully.

PRESENTATION

EXIFLEA® PLUS for	Pack colour	Contents	APVMA Approval No.
Puppies & small dogs up to 4kg	Green	0.4mL	86382/150409
Dogs 4-10kg	Turquoise	1.0mL	86383/150405
Dogs 10-25kg	Maroon	2.5mL	86384/150408
Dogs over 25kg	Blue	4.0mL	86385/150407

GENERAL DIRECTIONS

FLEA CONTROL AND PREVENTION

EXIFLEA® PLUS has been shown to kill adult and larval fleas within 20 minutes of contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite the animal to receive a lethal dose.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with EXIFLEA® PLUS rapidly kills fleas and reduces the incidence of this condition.

EXIFLEA® PLUS may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the dog. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product.

It is recommended that all dogs in the same environment should be protected by EXIFLEA® PLUS for dogs and all cats with EXIFLEA® PLUS for cats.

HEARTWORM PREVENTION

Treatment with EXIFLEA® PLUS should start one month before the first exposure of mosquitoes.

EXIFLEA® PLUS must be applied at one-month intervals during the time of the year when mosquitoes (which carry and transmit the heartworm larvae) are present. In many areas of Australia continuous, year-round heart worm prevention is recommended. To establish a routine, it is recommended that the same day or date be used each month to administer EXIFLEA® PLUS. When replacing another heartworm preventive product, the first treatment with EXIFLEA® PLUS must be given within one month of the last dose of the previous product.

WORM CONTROL

EXIFLEA® PLUS treats and controls major gastrointestinal nematodes of dogs but does not control tapeworms. Dogs can become reinfested with worms, but regular use of EXIFLEA® PLUS will remove them. If worm problems persist consult a veterinarian.

TREATMENT AND CONTROL OF GENERALISED DEMODICOSIS

EXIFLEA® PLUS has been shown to be highly effective in the treatment of generalised demodicosis when administered at monthly intervals for 2 to 4 treatments.

TREATMENT AND CONTROL OF SARCOPTIC MANGE

EXIFLEA® PLUS has been shown to eliminate 100% of sarcoptic mites (*Sarcoptes scabiei*) after a single dose. A second dose of EXIFLEA® PLUS a month later will result in continued improvement of the clinical signs associated with sarcoptic mange. Monthly application of EXIFLEA® PLUS will control any subsequent sarcoptic mite infestation.

TREATMENT AND CONTROL OF EAR MITES

For the treatment of ear mites (*Otodectes cynotis*), EXIFLEA® PLUS is highly efficacious as a single dose. Monthly application of EXIFLEA® PLUS will control any subsequent ear mite infection. To maximise efficacy thorough cleaning of ear canal is recommended with a product that is registered for that purpose or as directed by a veterinarian.

PUPPIES

Puppies may be treated with EXIFLEA® PLUS from 7 weeks of age.

For control of roundworms and hookworms, puppies require worming at 2, 4 and 8 weeks of age. When worming puppies prior to commencing EXIFLEA® PLUS use a specific worming agent.

Once treatment with EXIFLEA® PLUS every month is commenced worms will be continuously controlled.

PREGNANCY AND LACTATION

No primary embryo-toxic, teratogenic or reproductive toxic effects have been observed with Imidacloprid or Moxidectin in laboratory species. Evidence suggests that no adverse effects are to be expected in pregnant and lactating dogs. However, safety of EXIFLEA® PLUS has not been established during pregnancy and lactation.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes. If product in eyes, wash it out immediately with water. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

DISPOSAL

Dispose of used applicators by wrapping with paper and putting in garbage.

STORAGE

Store below 25°C (Air Conditioning).

EXIFLEA® is a registered trademark of Abbey Laboratories Pty Ltd.

Harmful if swallowed.

Harmful if inhaled.

May cause damage to organs.

Wash thoroughly after handling.

Do not eat, drink or smoke when using this product.

IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.

ABBEE LABORATORIES PTY LTD
DISTRIBUTED BY:
ABBEE ANIMAL HEALTH PTY LTD
UNIT 27 / 1 MAITLAND PLACE
NORWEST NSW 2153
INFO LINE 02 8088 0720
www.abbeylabs.com.au

